

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

קידום בריאות

שיעור התיעוד של מרכיבי BMI בקרב בני 20-64

56.7%	55.6%	56.8%	59.3%	63.9%	67.4%	64.6%	62.8%	52.2%	54.4%	52.1%	50.3%	76.4%	75.2%	74.0%	76.5%	20-24	זכר
75.7%	77.3%	76.8%	80.0%	60.6%	64.7%	65.3%	63.8%	73.6%	75.2%	73.7%	73.3%	73.0%	74.7%	72.5%	71.8%	25-34	
85.3%	84.8%	82.8%	85.8%	65.1%	68.9%	71.0%	70.5%	76.2%	78.5%	78.0%	77.7%	80.0%	81.2%	79.9%	80.8%	35-44	
93.0%	92.1%	90.3%	91.8%	73.5%	77.6%	78.7%	80.4%	84.6%	85.1%	84.8%	86.0%	87.9%	89.1%	89.3%	90.5%	45-54	
95.0%	95.0%	94.1%	95.2%	82.3%	85.9%	87.3%	88.7%	91.8%	91.8%	91.1%	90.5%	95.7%	96.6%	96.4%	96.8%	55-64	
73.7%	72.8%	75.5%	79.2%	73.6%	79.4%	81.4%	82.3%	69.0%	67.9%	67.1%	68.9%	82.8%	83.7%	84.0%	85.5%	20-24	נקבה
89.3%	90.8%	93.0%	95.3%	79.0%	84.1%	86.1%	87.8%	88.6%	90.0%	91.0%	92.6%	82.7%	85.3%	85.0%	84.2%	25-34	
93.0%	92.5%	93.6%	95.4%	76.4%	80.4%	82.3%	84.1%	88.6%	89.4%	89.9%	91.4%	85.4%	87.3%	88.0%	88.7%	35-44	
94.7%	94.7%	94.0%	95.9%	75.6%	81.1%	83.7%	85.4%	89.0%	90.2%	90.2%	90.2%	90.0%	92.4%	92.7%	94.0%	45-54	
95.9%	96.1%	95.5%	96.5%	82.9%	86.9%	88.8%	90.0%	93.9%	94.1%	93.7%	93.8%	96.8%	97.5%	97.7%	98.3%	55-64	

שיעור ההמצאות של תת משקל בקרב בני 20-64

6.5%	7.0%	6.2%	5.2%	6.6%	7.7%	7.1%	6.4%	5.7%	4.6%	5.5%	5.8%	9.2%	8.7%	6.0%	5.5%	20-24	זכר
2.7%	3.0%	3.3%	3.2%	2.7%	2.6%	3.5%	3.7%	3.1%	3.3%	3.3%	3.7%	2.7%	3.1%	3.5%	3.0%	25-34	
1.3%	1.5%	1.9%	1.9%	0.9%	1.2%	1.7%	1.5%	1.3%	1.7%	1.8%	2.2%	1.2%	1.6%	1.9%	1.6%	35-44	
0.7%	0.8%	1.2%	1.1%	0.3%	0.5%	0.8%	0.8%	0.9%	1.0%	0.9%	1.2%	0.6%	0.9%	1.1%	1.1%	45-54	
0.5%	0.6%	0.9%	0.9%	0.3%	0.5%	0.6%	1.0%	0.6%	0.6%	1.1%	1.2%	0.4%	0.8%	1.2%	1.4%	55-64	
10.9%	10.0%	7.9%	6.9%	11.8%	10.3%	8.7%	7.2%	10.6%	10.5%	9.5%	7.0%	12.8%	12.3%	11.3%	9.0%	20-24	נקבה
6.8%	5.5%	4.7%	3.4%	6.3%	5.2%	4.6%	3.5%	7.4%	5.8%	4.5%	3.8%	7.3%	6.6%	6.6%	4.3%	25-34	
4.9%	3.7%	3.0%	2.0%	3.9%	3.1%	2.7%	2.0%	4.4%	3.3%	2.5%	1.7%	5.1%	4.4%	3.5%	1.9%	35-44	
2.8%	2.2%	1.8%	1.1%	2.9%	2.0%	1.7%	1.1%	2.7%	2.0%	1.6%	1.0%	2.9%	2.5%	1.9%	0.9%	45-54	
1.7%	1.4%	1.2%	0.9%	1.7%	1.4%	1.3%	0.7%	1.8%	1.5%	1.1%	0.9%	1.7%	1.4%	1.4%	0.9%	55-64	

שיעור ההמצאות של עודף משקל בקרב בני 20-64

23.2%	24.5%	25.4%	26.2%	20.2%	21.9%	23.0%	24.8%	24.7%	24.3%	27.4%	26.7%	21.3%	21.4%	24.5%	22.5%	20-24	זכר
28.7%	31.9%	32.9%	32.7%	28.6%	31.3%	31.6%	32.1%	27.5%	31.4%	32.5%	32.9%	28.6%	30.8%	31.8%	31.7%	25-34	
38.5%	39.8%	38.8%	38.4%	38.3%	39.3%	38.4%	36.7%	36.1%	39.4%	39.2%	38.4%	37.4%	38.2%	38.2%	39.2%	35-44	
44.3%	43.4%	41.4%	39.9%	45.3%	43.9%	41.5%	39.5%	43.8%	43.1%	43.0%	41.4%	44.2%	42.9%	42.0%	42.3%	45-54	
48.7%	45.8%	43.2%	41.7%	48.4%	46.2%	43.0%	42.2%	46.8%	45.4%	42.6%	41.4%	48.1%	45.0%	43.0%	42.0%	55-64	
21.2%	23.1%	24.9%	26.6%	18.2%	20.3%	24.8%	27.3%	17.5%	21.1%	23.6%	27.8%	14.9%	18.9%	20.0%	20.9%	20-24	נקבה
24.8%	27.2%	30.6%	31.9%	27.0%	29.3%	30.3%	32.5%	24.7%	28.3%	30.4%	32.5%	21.7%	24.3%	25.6%	28.7%	25-34	
26.8%	29.8%	31.8%	33.9%	29.9%	30.4%	31.1%	34.1%	27.3%	29.5%	32.7%	33.7%	24.7%	27.2%	29.4%	33.4%	35-44	
29.1%	31.5%	32.9%	33.4%	28.1%	31.0%	32.3%	32.4%	30.2%	31.4%	33.0%	32.7%	27.6%	30.5%	32.6%	33.8%	45-54	
34.3%	35.5%	34.9%	31.4%	33.6%	35.2%	34.7%	31.8%	34.5%	35.2%	34.3%	33.3%	33.7%	35.0%	34.2%	31.4%	55-64	

שיעור ההמצאות של השמנת יתר בקרב בני 20-64

13.2%	14.9%	16.2%	17.2%	11.1%	15.6%	16.6%	16.1%	11.3%	13.3%	14.9%	16.8%	11.5%	14.5%	14.3%	13.0%	20-24	זכר
10.6%	15.7%	19.4%	23.0%	11.6%	15.3%	19.6%	23.1%	11.0%	15.8%	18.5%	21.5%	11.0%	14.5%	17.2%	19.3%	25-34	
15.3%	21.8%	27.3%	32.1%	16.1%	21.8%	26.8%	32.5%	16.8%	21.1%	25.5%	30.5%	14.0%	18.6%	22.8%	25.5%	35-44	
22.3%	28.5%	32.8%	37.2%	21.9%	28.9%	33.5%	38.3%	24.0%	29.4%	31.5%	35.1%	19.6%	25.0%	28.6%	30.4%	45-54	
24.7%	29.9%	34.0%	35.3%	24.8%	29.8%	33.5%	36.2%	26.3%	30.2%	31.4%	34.0%	24.3%	27.9%	28.7%	30.7%	55-64	
12.7%	15.5%	16.4%	18.8%	10.5%	15.2%	17.2%	17.2%	9.8%	15.2%	17.0%	20.0%	11.8%	13.0%	13.7%	14.8%	20-24	נקבה
15.4%	21.7%	24.1%	27.4%	15.0%	21.4%	25.5%	27.8%	17.0%	22.9%	24.8%	28.4%	12.4%	18.1%	21.0%	24.6%	25-34	
18.1%	23.9%	29.8%	36.9%	18.4%	24.6%	29.8%	35.4%	19.2%	25.5%	30.2%	38.4%	15.3%	21.0%	26.6%	33.6%	35-44	
19.1%	26.1%	34.3%	44.8%	19.4%	27.2%	34.1%	43.4%	21.8%	28.3%	36.6%	46.4%	19.3%	26.1%	34.4%	44.7%	45-54	
22.2%	29.7%	39.3%	51.5%	22.0%	30.4%	37.8%	48.4%	25.1%	31.4%	42.0%	48.5%	23.6%	30.6%	39.0%	51.4%	55-64	

שיעור תיעוד מצב עישון בקרב בני 16-74

58.1%	54.0%	53.0%	58.8%	66.2%	71.0%	73.8%	73.0%	74.5%	75.5%	76.2%	74.1%	98.5%	99.2%	99.2%	99.7%	16-24	זכר
62.6%	60.4%	57.9%	63.3%	73.2%	76.4%	77.4%	77.8%	74.5%	74.0%	74.8%	74.7%	98.6%	99.2%	99.3%	99.7%	25-34	
69.2%	65.8%	64.6%	68.5%	56.8%	62.5%	65.5%	64.4%	76.5%	75.9%	77.2%	77.9%	90.8%	93.0%	94.2%	94.7%	35-44	
77.1%	73.9%	74.5%	77.7%	60.2%	67.2%	70.9%	71.2%	82.0%	83.2%	82.7%	84.7%	57.3%	63.6%	67.1%	68.0%	45-54	
84.4%	81.9%	83.3%	85.1%	72.6%	78.2%	80.9%	81.9%	90.0%	89.6%	89.2%	89.5%	64.6%	72.4%	75.5%	76.7%	55-64	
93.9%	92.8%	91.8%	91.5%	87.0%	89.4%	90.6%	90.1%	95.0%	94.3%	92.7%	92.2%	73.8%	77.0%	79.1%	79.7%	65-74	
63.7%	59.6%	60.9%	69.0%	77.0%	81.8%	82.3%	81.1%	82.8%	83.5%	83.9%	82.6%	98.9%	99.3%	99.6%	99.8%	16-24	נקבה
83.6%	84.1%	87.0%	89.8%	91.9%	93.1%	93.7%	94.3%	89.3%	90.5%	91.9%	93.2%	99.4%	99.7%	99.8%	100.0%	25-34	
84.5%	80.3%	83.6%	87.8%	71.1%	73.1%	75.0%	74.2%	89.1%	88.6%	89.8%	91.1%	91.4%	92.8%	92.4%	90.4%	35-44	
81.1%	78.1%	80.7%	84.9%	63.5%	70.0%	72.7%	72.2%	88.5%	88.9%	88.6%	90.0%	51.5%	53.6%	47.6%	33.6%	45-54	
86.0%	84.0%	85.9%	88.4%	73.7%	77.4%	80.5%	80.4%	92.8%	92.6%	91.6%	92.8%	59.7%	60.4%	57.8%	46.4%	55-64	
94.2%	92.4%	92.1%	92.4%	85.5%	87.8%	89.1%	88.8%	95.9%	94.9%	93.4%	94.2%	66.5%	65.4%	63.0%	53.2%	65-74	

שיעור עישון בקרב בני 16-74

מאוחדת				מכבי				לאומית				כללית				SES גיל	
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1		
10.1%	15.2%	20.8%	32.5%	6.4%	11.1%	14.7%	15.9%	21.7%	24.0%	29.7%	31.2%	6.8%	10.9%	14.8%	15.4%	16-24	זכר
31.6%	37.3%	40.8%	50.7%	21.6%	26.8%	30.3%	27.0%	36.5%	43.0%	49.3%	49.7%	23.6%	30.6%	35.7%	38.1%	25-34	
26.2%	34.0%	35.6%	45.0%	20.1%	27.5%	33.2%	30.8%	35.3%	39.8%	45.4%	49.4%	21.0%	29.8%	36.4%	41.4%	35-44	
20.6%	30.1%	35.5%	43.8%	15.9%	24.2%	32.3%	30.3%	27.7%	38.4%	44.2%	47.1%	26.1%	37.9%	48.1%	55.1%	45-54	
19.6%	28.5%	33.7%	42.1%	13.0%	21.6%	29.8%	29.9%	26.4%	34.0%	41.1%	43.0%	20.6%	32.8%	42.8%	50.6%	55-64	
14.5%	20.1%	23.5%	30.2%	9.4%	14.9%	19.3%	20.3%	20.0%	24.9%	31.4%	34.4%	12.6%	20.0%	28.4%	37.5%	65-74	נקבה
7.5%	7.6%	4.3%	4.8%	5.5%	6.6%	5.1%	1.7%	11.3%	12.6%	8.0%	5.4%	4.7%	5.5%	4.0%	1.2%	16-24	
19.6%	18.8%	9.4%	7.3%	10.6%	12.4%	11.8%	5.2%	24.2%	24.1%	15.5%	10.6%	13.0%	15.3%	12.7%	4.5%	25-34	
17.1%	20.1%	12.3%	8.2%	10.3%	15.0%	16.8%	8.3%	23.7%	25.0%	17.1%	11.6%	13.0%	17.4%	15.7%	6.8%	35-44	
15.9%	20.0%	15.9%	10.7%	11.8%	16.8%	19.3%	11.0%	21.8%	25.4%	19.0%	13.1%	21.1%	29.6%	30.7%	22.1%	45-54	
16.1%	20.6%	17.5%	12.8%	11.9%	16.3%	19.0%	12.5%	21.4%	24.6%	20.9%	17.5%	19.8%	27.4%	29.0%	24.7%	55-64	
12.1%	14.6%	13.1%	9.4%	9.1%	11.3%	11.6%	8.6%	16.8%	19.1%	16.4%	14.0%	12.6%	17.5%	19.5%	17.8%	65-74	

מאחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

בדיקות סקר לגילוי מוקדם של סרטן

שיעור הביצוע של ממוגרפיה לגילוי מוקדם של סרטן השד בנשים בנות 74-50

78.6%	74.9%	68.3%	61.9%	81.8%	80.5%	76.7%	71.7%	68.5%	64.4%	58.4%	52.9%	75.5%	71.7%	68.3%	67.9%	50-54	נקבה
75.0%	74.3%	68.0%	63.2%	81.5%	80.5%	76.9%	73.0%	63.9%	62.8%	57.6%	51.2%	73.4%	70.4%	67.8%	67.4%	55-59	
74.3%	73.7%	69.6%	63.8%	81.4%	80.5%	78.8%	74.5%	66.1%	65.0%	59.0%	52.7%	74.4%	71.6%	70.2%	68.6%	60-64	
75.6%	73.4%	70.0%	61.8%	81.3%	81.0%	79.4%	75.6%	66.3%	65.7%	59.3%	55.0%	76.3%	73.5%	69.6%	68.2%	65-69	
73.2%	72.8%	65.8%	58.1%	77.2%	75.6%	73.7%	69.8%	62.8%	64.2%	57.0%	51.5%	72.3%	67.9%	62.0%	54.2%	70-74	

שיעור הביצוע של בדיקה לגילוי מוקדם של סרטן המעי הגס בקרב בני 74-50

56.9%	51.9%	46.8%	44.1%	63.7%	59.1%	53.9%	51.8%	50.3%	44.2%	39.4%	39.5%	55.4%	49.4%	45.4%	46.5%	50-54	זכר
65.7%	61.1%	56.1%	53.9%	73.7%	69.0%	64.0%	61.6%	59.0%	53.6%	49.8%	46.0%	65.1%	57.7%	53.1%	53.7%	55-59	
71.0%	67.7%	62.4%	58.5%	77.0%	73.5%	68.9%	67.9%	64.8%	61.2%	57.1%	51.6%	68.3%	63.5%	58.2%	59.1%	60-64	
75.0%	72.4%	66.3%	63.8%	79.4%	77.3%	75.3%	72.6%	66.6%	67.9%	60.8%	58.1%	74.0%	68.6%	64.0%	63.9%	65-69	
77.8%	74.6%	70.4%	66.7%	81.4%	79.5%	76.8%	71.8%	73.7%	70.2%	65.6%	62.3%	75.8%	70.9%	65.1%	64.4%	70-74	
55.5%	53.3%	51.7%	50.9%	61.9%	60.7%	58.0%	56.5%	49.8%	49.9%	46.9%	44.8%	54.8%	51.2%	51.4%	55.1%	50-54	נקבה
67.5%	62.9%	58.1%	56.4%	72.0%	69.8%	66.3%	65.2%	60.5%	57.9%	54.4%	50.6%	64.1%	59.3%	57.7%	60.7%	55-59	
69.9%	67.4%	64.4%	61.6%	74.9%	73.7%	72.3%	69.2%	67.1%	64.8%	60.1%	55.3%	68.3%	64.6%	62.0%	64.3%	60-64	
73.6%	73.2%	69.5%	66.3%	78.6%	77.2%	75.6%	74.2%	71.8%	69.0%	65.5%	59.4%	72.5%	69.3%	65.0%	66.4%	65-69	
74.8%	74.7%	69.4%	64.7%	78.4%	77.5%	75.9%	73.5%	71.0%	71.1%	66.5%	64.8%	72.8%	69.6%	64.4%	61.8%	70-74	

שיעור המבוטחים בעלי בדיקת דם סמוי חיובית, אשר עברו בדיקת קולונסקופיה תוך 180 ימים מיום ביצוע בדיקת הדם הסמוי

80.5%	57.7%	60.6%	55.6%	81.5%	77.5%	73.8%	73.1%	-	64.9%	62.0%	42.9%	75.4%	70.0%	60.1%	56.6%	50-54	זכר
59.1%	60.2%	64.1%	45.0%	78.9%	81.8%	71.3%	76.1%	-	45.0%	43.7%	37.0%	72.4%	62.3%	57.4%	51.3%	55-59	
78.7%	61.7%	56.1%	54.6%	81.2%	76.1%	66.5%	64.4%	-	61.8%	48.7%	46.5%	74.6%	70.4%	58.3%	52.4%	60-64	
63.3%	59.8%	52.6%	50.7%	70.6%	76.3%	66.2%	67.5%	-	51.7%	44.4%	44.7%	78.2%	67.3%	60.7%	51.3%	65-69	
59.2%	54.1%	56.8%	56.9%	77.1%	73.2%	65.9%	55.0%	-	52.3%	43.0%	45.2%	74.0%	70.0%	59.4%	49.2%	70-74	
-	62.4%	62.0%	61.4%	88.2%	82.4%	76.9%	67.7%	-	75.8%	50.0%	54.8%	78.9%	71.3%	62.3%	55.3%	50-54	נקבה
63.9%	61.8%	55.3%	58.4%	87.4%	82.2%	83.4%	72.6%	-	-	55.6%	52.5%	78.5%	75.2%	66.7%	59.4%	55-59	
61.1%	59.3%	55.9%	57.0%	83.2%	78.4%	78.7%	84.0%	-	60.4%	57.8%	54.1%	72.2%	74.2%	63.4%	52.2%	60-64	
65.6%	59.6%	60.7%	56.6%	71.1%	79.8%	68.8%	73.0%	-	56.1%	52.8%	41.9%	78.0%	72.5%	60.0%	50.6%	65-69	
67.2%	58.1%	54.3%	44.4%	74.3%	79.1%	72.1%	70.8%	-	62.7%	47.6%	45.2%	76.0%	70.9%	59.6%	48.6%	70-74	

שיעור הנשים בגילאים 54-25 אשר ביצעו בדיקות סיקור לסרטן צוואר הרחם ב-3 השנים האחרונות

59.5%	53.4%	41.4%	33.3%	67.9%	65.7%	59.7%	46.2%	60.8%	54.9%	45.5%	35.8%	64.6%	59.9%	49.7%	35.7%	25-29	נקבה
65.7%	60.3%	49.6%	42.5%	72.0%	69.7%	62.8%	49.9%	64.4%	59.7%	52.1%	42.6%	70.1%	64.6%	54.9%	42.1%	30-34	
64.7%	59.2%	49.7%	41.7%	70.0%	67.6%	61.3%	50.4%	64.9%	61.0%	51.3%	42.4%	67.1%	62.4%	52.2%	40.3%	35-39	
63.2%	59.3%	49.1%	40.6%	67.5%	64.7%	58.8%	49.7%	62.4%	58.8%	50.4%	40.3%	64.5%	59.8%	49.8%	38.5%	40-44	
64.4%	59.0%	48.8%	38.5%	66.9%	63.4%	57.7%	49.6%	60.7%	57.2%	49.2%	39.8%	62.2%	56.8%	46.2%	36.4%	45-49	
59.6%	53.8%	44.1%	34.4%	66.6%	62.9%	56.0%	47.7%	56.9%	52.4%	43.5%	36.3%	57.5%	50.3%	39.5%	30.7%	50-54	

שיעור הנשים בגילאים 54-30 אשר לא ביצעו בדיקת סיקור לסרטן צוואר הרחם ב-5 השנים האחרונות

20.7%	26.7%	38.3%	48.6%	11.5%	12.6%	18.2%	30.6%	24.3%	26.6%	35.0%	46.6%	16.3%	21.4%	32.0%	48.3%	30-34	נקבה
19.3%	24.4%	35.9%	46.6%	11.5%	12.7%	17.8%	28.9%	22.1%	23.9%	32.7%	44.7%	17.0%	21.4%	32.0%	46.9%	35-39	
20.7%	24.9%	36.2%	47.8%	13.1%	14.1%	19.6%	27.5%	23.2%	25.6%	34.6%	45.5%	19.7%	24.0%	34.4%	48.4%	40-44	
20.6%	25.2%	36.8%	49.4%	14.6%	16.3%	21.5%	27.4%	25.2%	26.3%	35.0%	46.4%	23.0%	27.6%	38.5%	50.3%	45-49	
23.5%	29.1%	39.6%	51.6%	16.5%	17.8%	23.3%	29.9%	28.8%	29.7%	38.8%	48.0%	26.7%	32.8%	43.7%	55.5%	50-54	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

ילדים ובני נוער

שיעור הביצוע של בדיקת המוגלובין בתינוקות בני 9-18 חודשים

86.2%	84.0%	84.0%	85.5%	84.9%	84.6%	78.1%	77.8%	80.2%	75.7%	69.1%	68.0%	87.8%	87.1%	85.1%	87.5%	1	זכר
86.8%	84.4%	84.6%	84.5%	85.1%	83.2%	77.8%	77.2%	76.5%	71.6%	69.9%	67.3%	88.3%	86.6%	84.3%	87.2%	1	נקבה

שיעור ההימצאות של אנמיה בקרב תינוקות בני 9-18 חודשים

9.4%	8.1%	10.5%	13.1%	8.9%	8.8%	9.6%	13.7%	8.6%	8.0%	9.6%	13.4%	9.6%	9.0%	9.7%	9.2%	1	זכר
7.0%	8.0%	9.1%	10.4%	8.9%	8.3%	9.2%	11.7%	6.5%	7.1%	8.7%	11.1%	8.3%	8.5%	8.9%	9.0%	1	נקבה

שיעור התייעוד של מדידת גובה ומשקל בקרב ילדים בני 7

67.1%	65.9%	64.7%	64.2%	51.7%	50.1%	47.6%	43.5%	65.9%	69.3%	61.4%	55.4%	79.0%	77.4%	75.6%	75.9%	7	זכר
69.5%	63.6%	63.3%	63.1%	49.0%	49.0%	45.2%	41.6%	66.6%	69.2%	57.6%	52.4%	78.6%	77.0%	74.4%	73.9%	7	נקבה

שיעור ההימצאות של רזון (Thinness) בקרב ילדים שמלאו להם 7 שנים בשנת המדד

4.9%	4.7%	5.8%	5.8%	3.1%	3.5%	3.6%	3.5%	4.6%	5.6%	4.6%	2.0%	3.8%	4.0%	4.9%	6.6%	7	זכר
4.2%	3.8%	4.9%	4.6%	2.0%	2.2%	2.0%	1.7%	3.2%	3.3%	2.7%	3.3%	2.3%	3.0%	3.9%	5.9%	7	נקבה

שיעור ההימצאות של משקל תקין בקרב ילדים שמלאו להם 7 שנים בשנת המדד

75.5%	73.0%	72.4%	73.6%	78.6%	74.5%	73.5%	75.7%	75.1%	73.1%	76.3%	76.3%	78.8%	75.6%	72.8%	74.3%	7	זכר
76.3%	71.8%	73.4%	74.6%	76.6%	74.5%	73.8%	77.5%	72.5%	74.2%	75.7%	76.7%	79.2%	75.8%	73.4%	76.5%	7	נקבה

שיעור ההימצאות של עודף משקל בקרב ילדים שמלאו להם 7 שנים בשנת המדד

13.2%	13.2%	13.4%	11.6%	11.1%	12.5%	12.3%	12.8%	12.3%	12.4%	10.1%	12.5%	11.2%	11.6%	12.3%	10.1%	7	זכר
12.8%	16.1%	13.9%	12.7%	13.6%	13.6%	13.1%	10.5%	12.1%	13.4%	12.1%	11.7%	12.5%	12.8%	13.3%	9.8%	7	נקבה

שיעור ההימצאות של השמנת יתר בקרב ילדים שמלאו להם 7 שנים בשנת המדד

6.4%	9.1%	8.4%	9.0%	7.1%	9.5%	10.6%	8.0%	8.0%	9.0%	9.1%	9.2%	6.2%	8.8%	10.1%	9.0%	7	זכר
6.8%	8.2%	7.8%	8.1%	7.8%	9.8%	11.2%	10.3%	12.1%	9.2%	9.5%	8.3%	6.1%	8.4%	9.4%	7.8%	7	נקבה

שיעור התייעוד של מרכיבי BMI בקרב בני 14-15

80.8%	76.4%	74.1%	75.2%	62.3%	62.3%	58.8%	54.6%	82.0%	79.4%	74.5%	65.5%	78.7%	78.4%	75.1%	76.3%	14	זכר
86.9%	79.6%	76.5%	76.9%	67.6%	65.9%	62.0%	53.1%	70.5%	79.3%	74.4%	65.6%	75.4%	72.8%	64.6%	51.9%	15	זכר
80.1%	73.2%	69.9%	76.6%	59.6%	61.0%	57.2%	51.8%	77.3%	76.8%	69.9%	62.3%	78.3%	79.3%	75.2%	75.3%	14	נקבה
84.5%	79.6%	74.5%	79.4%	67.1%	66.2%	60.3%	53.3%	76.3%	77.2%	75.0%	56.0%	75.8%	72.6%	65.2%	53.4%	15	נקבה

שיעור ההימצאות של רזון (Thinness) בקרב בני 14-15

4.0%	4.3%	4.0%	3.9%	4.7%	5.0%	4.7%	5.1%	3.9%	4.5%	4.5%	4.7%	4.5%	4.7%	4.9%	4.3%	14	זכר
3.2%	3.1%	4.0%	4.0%	3.6%	4.2%	4.8%	3.8%	9.7%	2.3%	4.8%	6.5%	3.7%	4.4%	4.8%	3.8%	15	זכר
2.6%	2.8%	3.8%	2.5%	2.4%	3.1%	3.7%	3.6%	4.1%	2.9%	3.5%	3.3%	2.8%	2.8%	3.1%	2.6%	14	נקבה
2.2%	2.0%	3.0%	2.2%	2.0%	2.6%	2.5%	3.6%	-	0.0%	3.8%	3.2%	2.2%	2.7%	2.6%	2.5%	15	נקבה

שיעור ההימצאות של משקל תקין בקרב בני 14-15

68.8%	63.7%	63.7%	59.1%	69.7%	63.7%	59.3%	56.1%	64.9%	61.4%	60.9%	60.1%	69.3%	64.2%	59.8%	60.2%	14	זכר
69.4%	66.5%	64.6%	58.1%	72.2%	63.9%	62.3%	61.1%	71.0%	60.2%	60.8%	56.1%	70.8%	64.7%	59.9%	57.0%	15	זכר
72.3%	66.6%	65.8%	64.2%	71.7%	65.7%	61.7%	63.8%	66.1%	63.9%	65.1%	65.0%	73.0%	67.1%	63.1%	65.5%	14	נקבה
74.1%	70.2%	69.6%	68.3%	73.8%	68.2%	65.3%	65.5%	-	53.8%	65.2%	72.0%	72.6%	68.3%	65.6%	63.9%	15	נקבה

שיעור ההימצאות של עודף משקל בקרב בני 14-15

18.2%	18.8%	19.2%	21.0%	16.3%	18.0%	18.7%	21.1%	18.0%	18.8%	18.9%	17.3%	17.2%	17.7%	19.0%	18.5%	14	זכר
17.5%	18.5%	17.5%	21.5%	15.8%	18.7%	17.7%	17.7%	9.7%	18.2%	21.6%	18.7%	16.6%	18.0%	18.3%	20.1%	15	זכר
18.0%	19.2%	20.1%	21.8%	17.5%	19.0%	20.4%	19.1%	19.7%	18.6%	18.6%	18.6%	16.2%	18.0%	19.4%	19.5%	14	נקבה
16.7%	17.0%	18.1%	19.7%	16.7%	17.8%	19.7%	19.9%	-	28.2%	20.5%	14.0%	16.9%	17.4%	18.5%	19.1%	15	נקבה

שיעור ההימצאות של השמנת יתר בקרב בני 14-15

9.1%	13.3%	13.1%	16.0%	9.3%	13.3%	17.3%	17.8%	13.2%	15.3%	15.7%	18.0%	9.1%	13.4%	16.2%	17.0%	14	זכר
10.0%	11.9%	13.9%	16.4%	8.4%	13.1%	15.2%	17.4%	9.7%	19.3%	12.8%	18.7%	8.8%	12.9%	17.0%	19.1%	15	זכר
7.1%	11.3%	10.3%	11.5%	8.4%	12.2%	14.2%	13.5%	10.1%	14.6%	12.8%	13.0%	8.0%	12.1%	14.4%	12.3%	14	נקבה
7.0%	10.7%	9.4%	9.8%	7.4%	11.4%	12.6%	10.9%	-	17.9%	10.6%	10.8%	8.3%	11.5%	13.2%	14.5%	15	נקבה

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

מבוגרים בני 65 ומעלה

שיעור קבלת חיסון נגד שפעת בקרב בני 65 ומעלה

59.9%	50.6%	42.0%	43.2%	58.7%	48.1%	40.3%	37.6%	71.9%	67.4%	64.0%	61.8%	63.3%	55.7%	50.7%	52.6%	65-69	זכר
71.7%	62.1%	51.8%	49.9%	69.1%	57.8%	48.9%	45.3%	79.8%	73.8%	70.5%	70.5%	75.0%	67.2%	60.0%	59.6%	70-74	
77.9%	68.9%	57.8%	53.9%	75.0%	63.8%	54.3%	50.6%	86.2%	79.7%	72.2%	72.0%	79.8%	73.6%	65.3%	65.1%	75-79	
77.7%	70.0%	60.7%	58.6%	74.6%	64.1%	53.5%	49.9%	83.5%	75.3%	73.1%	73.6%	82.1%	74.8%	67.5%	66.4%	80-84	
77.2%	70.4%	61.3%	63.1%	74.4%	65.0%	57.1%	54.0%	84.5%	77.3%	70.8%	66.1%	81.3%	76.1%	69.4%	66.2%	85+	
59.4%	47.7%	39.4%	38.2%	58.0%	46.2%	36.9%	33.8%	73.7%	69.6%	67.9%	66.9%	61.8%	52.5%	46.1%	50.2%	65-69	נקבה
67.7%	57.9%	47.3%	43.3%	66.9%	53.6%	43.9%	40.3%	80.8%	76.4%	70.6%	71.7%	71.4%	62.2%	54.5%	53.8%	70-74	
73.6%	61.7%	52.0%	47.6%	70.4%	56.8%	48.2%	45.8%	82.3%	77.3%	72.9%	71.9%	75.3%	67.4%	59.4%	58.6%	75-79	
71.0%	61.2%	51.6%	50.3%	68.0%	55.7%	46.3%	45.3%	76.5%	74.3%	69.6%	71.6%	75.1%	66.9%	61.2%	60.2%	80-84	
66.5%	62.8%	55.5%	53.6%	67.1%	54.5%	47.4%	46.6%	76.3%	69.9%	68.0%	68.0%	73.4%	68.3%	63.6%	60.7%	85+	

שיעור קבלת חיסון נגד פניאומוקוק בקרב בני 65-74

73.6%	69.5%	64.6%	69.0%	74.6%	71.6%	69.5%	70.3%	90.4%	90.0%	87.4%	87.4%	72.1%	69.1%	68.3%	73.5%	65-69	זכר
89.7%	85.4%	82.9%	85.1%	89.5%	87.8%	87.7%	87.1%	97.7%	97.3%	96.8%	96.2%	88.0%	86.1%	84.9%	87.8%	70-74	
72.8%	65.8%	61.0%	65.5%	72.1%	68.9%	65.8%	65.4%	93.6%	91.5%	90.6%	90.2%	71.1%	66.5%	64.3%	70.9%	65-69	נקבה
85.3%	81.8%	77.8%	79.7%	86.0%	83.9%	82.1%	83.1%	98.4%	98.0%	96.8%	96.5%	85.3%	82.0%	80.7%	84.8%	70-74	

שיעור התייעוד של מרכיבי BMI בקרב בני 65-84

71.7%	71.7%	70.4%	75.1%	52.4%	56.9%	59.4%	62.8%	66.5%	66.8%	66.8%	66.0%	70.9%	73.0%	72.6%	74.4%	65-69	זכר
79.2%	78.2%	75.8%	77.3%	56.8%	62.9%	64.2%	66.5%	77.6%	74.9%	73.1%	72.0%	80.8%	81.9%	80.6%	82.0%	70-74	
79.7%	80.0%	77.6%	78.2%	60.0%	65.9%	65.6%	70.4%	54.0%	56.5%	53.1%	52.5%	82.9%	84.3%	81.3%	82.9%	75-79	
52.8%	50.5%	49.0%	48.4%	62.2%	66.0%	66.0%	67.5%	52.2%	49.2%	46.8%	48.2%	84.1%	84.9%	82.6%	81.4%	80-84	
73.0%	72.6%	73.1%	77.4%	52.1%	57.4%	59.7%	64.3%	71.7%	68.3%	69.4%	71.6%	71.3%	73.7%	74.7%	78.2%	65-69	נקבה
77.9%	79.1%	76.5%	78.9%	56.2%	61.1%	63.6%	66.5%	77.8%	77.5%	74.9%	75.9%	80.9%	82.0%	81.8%	83.0%	70-74	
79.1%	78.4%	75.3%	78.0%	56.9%	63.0%	64.4%	69.4%	53.2%	54.2%	53.1%	58.3%	82.0%	83.3%	81.9%	82.8%	75-79	
48.2%	46.2%	43.6%	48.1%	56.4%	60.9%	62.1%	67.2%	46.5%	46.7%	42.5%	46.6%	79.9%	81.0%	80.1%	79.2%	80-84	

שיעור המבוגרים בני 65 ומעלה שחלה ירידה חדה במשקלם בשנתיים האחרונות

5.5%	6.2%	6.0%	6.9%	7.6%	7.9%	8.3%	8.4%	6.2%	7.4%	6.7%	6.8%	5.5%	5.7%	5.8%	5.7%	65-69	זכר
6.1%	6.4%	6.9%	8.5%	8.0%	8.1%	8.7%	8.1%	7.1%	6.7%	7.6%	6.5%	5.5%	6.1%	5.8%	6.3%	70-74	
6.9%	6.9%	7.2%	7.5%	9.8%	9.4%	9.9%	9.4%	8.5%	10.0%	9.4%	9.7%	5.5%	6.5%	7.0%	6.5%	75-79	
7.8%	7.3%	8.9%	8.7%	9.7%	10.3%	9.7%	10.8%	9.4%	10.4%	11.4%	9.3%	7.4%	7.3%	7.6%	8.0%	80-84	
7.8%	9.5%	9.6%	12.0%	12.4%	13.7%	15.5%	13.6%	10.8%	10.9%	11.4%	7.9%	7.6%	8.3%	8.6%	8.4%	85+	
7.1%	8.1%	8.8%	9.0%	9.3%	11.3%	11.2%	10.5%	7.0%	8.7%	8.9%	8.4%	7.4%	7.8%	8.1%	6.6%	65-69	נקבה
7.6%	8.6%	8.8%	8.1%	10.5%	10.5%	11.0%	10.4%	10.2%	8.4%	8.9%	7.8%	6.9%	8.1%	7.6%	7.0%	70-74	
8.1%	8.5%	8.5%	10.4%	10.3%	11.0%	10.7%	12.1%	9.4%	11.1%	10.1%	8.6%	7.3%	7.6%	8.3%	7.2%	75-79	
7.7%	10.4%	12.4%	8.7%	11.4%	11.6%	12.3%	12.4%	9.9%	10.0%	9.7%	12.3%	7.8%	8.3%	8.3%	8.4%	80-84	
10.5%	12.2%	11.0%	14.2%	16.9%	15.1%	14.5%	18.1%	11.1%	8.7%	12.2%	10.1%	8.6%	9.8%	9.6%	9.5%	85+	

שיעור תת משקל בקרב מבוגרים בני 75 ומעלה

11.4%	11.2%	10.6%	13.1%	12.0%	11.1%	12.4%	13.0%	10.8%	13.5%	11.6%	17.7%	14.0%	13.4%	14.4%	15.2%	75-79	זכר
19.2%	13.4%	11.6%	14.8%	14.3%	13.5%	14.0%	14.9%	14.2%	14.6%	15.2%	16.2%	16.5%	15.7%	16.5%	17.5%	80-84	
18.4%	15.3%	16.7%	17.4%	20.5%	18.4%	18.2%	18.9%	19.5%	23.1%	17.1%	23.8%	19.5%	20.2%	21.1%	21.4%	85+	
22.5%	16.1%	12.8%	10.0%	20.7%	15.9%	11.9%	9.5%	16.8%	15.5%	11.6%	11.9%	20.3%	16.5%	13.7%	9.8%	75-79	נקבה
21.5%	18.1%	13.6%	13.0%	22.1%	14.4%	12.6%	12.4%	21.4%	18.4%	12.7%	13.8%	21.0%	17.9%	15.0%	12.5%	80-84	
26.2%	20.2%	16.0%	21.1%	26.8%	19.7%	16.6%	16.8%	20.3%	20.1%	15.8%	20.2%	23.9%	21.5%	19.3%	16.3%	85+	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור ההימצאות של השמנת יתר בקרב בני 65-84

27.4%	31.7%	33.5%	35.8%	26.1%	31.7%	33.6%	36.5%	25.9%	30.2%	32.5%	32.2%	25.7%	28.1%	29.4%	31.4%	65-69	זכר
24.6%	29.9%	33.7%	34.6%	24.8%	29.6%	33.0%	36.6%	24.3%	30.7%	31.6%	32.2%	23.9%	26.8%	28.0%	30.0%	70-74	
23.6%	26.2%	31.5%	33.3%	21.0%	26.5%	30.4%	29.2%	24.9%	26.1%	27.2%	30.3%	20.2%	23.3%	26.3%	28.7%	75-79	
19.7%	27.7%	25.9%	27.9%	19.2%	23.0%	24.6%	27.5%	17.5%	21.2%	25.2%	29.0%	17.3%	20.4%	22.1%	24.0%	80-84	
24.6%	33.4%	42.8%	55.6%	25.5%	35.2%	43.6%	50.7%	24.9%	34.1%	43.2%	51.3%	26.8%	33.6%	41.7%	53.6%	65-69	נקבה
25.3%	34.9%	42.7%	51.5%	26.5%	36.2%	44.4%	50.8%	25.5%	34.8%	44.3%	50.5%	26.4%	32.5%	41.0%	51.0%	70-74	
24.3%	33.8%	40.7%	48.4%	24.4%	33.6%	42.0%	49.8%	29.5%	33.4%	43.0%	49.0%	24.9%	31.7%	38.7%	49.6%	75-79	
23.5%	32.7%	40.3%	42.6%	24.7%	33.0%	41.0%	43.6%	26.3%	31.9%	38.8%	40.2%	24.3%	30.4%	36.6%	43.3%	80-84	

שיעור שימוש היתר בבנזודיאזפינים בקרב בני 65 ומעלה

1.4%	1.6%	2.0%	1.7%	1.5%	1.9%	2.0%	2.1%	1.6%	2.3%	2.2%	2.8%	2.0%	2.4%	2.6%	1.9%	65-69	זכר
1.8%	2.0%	3.1%	1.9%	2.2%	2.1%	2.6%	1.8%	3.2%	4.0%	3.6%	3.6%	3.3%	3.3%	3.3%	2.6%	70-74	
3.1%	2.6%	3.4%	2.0%	2.8%	2.7%	3.0%	2.6%	4.9%	5.1%	3.9%	3.7%	4.9%	4.9%	4.3%	3.5%	75-79	
3.4%	5.0%	4.9%	4.3%	3.8%	3.6%	4.1%	4.1%	8.0%	7.6%	8.5%	6.8%	8.1%	7.1%	6.4%	4.6%	80-84	
5.1%	5.8%	6.5%	5.0%	4.6%	4.5%	5.2%	5.5%	6.9%	8.7%	10.3%	10.6%	10.7%	10.3%	9.6%	6.3%	85+	
1.6%	2.1%	3.0%	2.0%	2.3%	2.5%	3.1%	2.6%	2.8%	3.5%	4.0%	3.2%	3.1%	3.2%	3.1%	2.1%	65-69	נקבה
2.8%	3.6%	4.1%	3.6%	3.2%	3.2%	3.5%	3.8%	5.7%	4.5%	5.3%	3.7%	4.9%	4.7%	4.3%	3.3%	70-74	
4.2%	4.1%	4.4%	3.1%	4.1%	4.5%	4.1%	4.3%	7.0%	7.2%	5.9%	5.6%	7.1%	6.5%	5.8%	3.6%	75-79	
5.6%	6.5%	7.1%	6.1%	4.9%	5.8%	6.4%	5.0%	9.4%	10.0%	9.7%	9.8%	10.5%	10.1%	8.7%	6.0%	80-84	
7.7%	8.4%	9.3%	7.3%	6.6%	6.3%	6.8%	5.8%	14.8%	11.4%	12.9%	11.6%	15.2%	14.2%	12.7%	8.9%	85+	

שיעור השימוש בבנזודיאזפינים ארוכי טווח בקרב בני 65 ומעלה

1.4%	1.3%	1.3%	2.2%	1.1%	1.0%	1.4%	1.5%	1.5%	1.0%	1.2%	2.5%	1.3%	1.2%	1.6%	1.6%	65-69	זכר
1.7%	1.3%	1.3%	2.0%	1.5%	1.3%	1.2%	1.8%	1.6%	1.1%	1.4%	1.1%	1.4%	1.3%	1.5%	1.6%	70-74	
1.9%	1.8%	1.4%	1.8%	1.8%	1.5%	1.2%	1.2%	0.8%	1.5%	1.7%	1.3%	1.5%	1.4%	1.5%	1.5%	75-79	
1.9%	2.0%	1.2%	2.4%	1.9%	1.4%	1.2%	1.1%	1.7%	1.6%	1.6%	1.8%	1.4%	1.6%	1.5%	1.2%	80-84	
1.3%	1.6%	1.2%	2.1%	1.8%	1.3%	1.0%	0.8%	1.2%	1.2%	1.5%	1.8%	1.8%	1.6%	1.3%	1.3%	85+	
1.9%	1.6%	1.9%	2.3%	1.8%	1.6%	1.4%	1.8%	1.7%	1.0%	1.5%	1.0%	1.5%	1.5%	1.6%	1.3%	65-69	נקבה
2.1%	2.2%	1.7%	2.1%	2.3%	1.7%	1.4%	1.8%	2.2%	1.4%	1.6%	1.3%	2.1%	1.8%	1.7%	1.4%	70-74	
2.4%	2.5%	2.4%	2.6%	2.5%	2.4%	1.7%	2.4%	2.9%	1.5%	1.8%	2.1%	2.3%	2.0%	1.7%	1.5%	75-79	
2.1%	2.0%	1.9%	1.8%	2.2%	2.2%	1.7%	2.3%	1.7%	1.5%	1.6%	1.1%	2.7%	2.3%	1.8%	1.2%	80-84	
2.5%	1.7%	2.5%	1.9%	2.7%	1.5%	1.9%	2.0%	2.2%	1.8%	2.1%	1.9%	2.4%	2.0%	1.7%	1.4%	85+	

מאחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

מחלות נשימתיות

שיעור ההימצאות של אסתמה מתמשכת בקרב בני 44-5

1.5%	1.2%	0.8%	0.6%	1.3%	1.3%	1.0%	0.8%	1.9%	1.3%	1.1%	0.6%	1.0%	1.0%	0.9%	0.7%	5-14	זכר
0.9%	1.0%	0.6%	0.4%	0.9%	0.7%	0.5%	0.3%	1.1%	1.1%	0.6%	0.4%	0.7%	0.6%	0.5%	0.4%	15-24	
1.2%	1.2%	0.7%	0.6%	0.8%	0.8%	0.7%	0.6%	1.6%	1.2%	0.8%	0.5%	0.7%	0.7%	0.6%	0.5%	25-34	
1.7%	1.2%	1.0%	1.0%	1.1%	1.0%	1.1%	0.7%	1.7%	1.5%	1.2%	1.1%	0.9%	1.0%	0.8%	0.9%	35-44	
0.7%	0.7%	0.5%	0.3%	0.8%	0.8%	0.7%	0.5%	1.3%	1.0%	0.4%	0.3%	0.6%	0.6%	0.5%	0.4%	5-14	נקבה
1.1%	0.7%	0.4%	0.3%	0.7%	0.6%	0.5%	0.3%	1.3%	0.9%	0.6%	0.3%	0.6%	0.5%	0.4%	0.3%	15-24	
1.2%	1.0%	0.6%	0.4%	0.9%	0.8%	0.7%	0.5%	1.4%	1.3%	0.8%	0.7%	0.7%	0.7%	0.7%	0.5%	25-34	
1.3%	1.4%	1.2%	0.9%	1.1%	1.3%	1.3%	0.9%	1.8%	1.8%	1.3%	1.2%	0.9%	1.0%	1.0%	1.0%	35-44	

שיעור יחס התרופות לאסתמה (AMR) גדול או שווה 0.5 בקרב בני 44-5

81.2%	80.7%	82.1%	83.2%	75.1%	73.7%	73.5%	77.0%	84.0%	76.3%	73.7%	76.4%	74.8%	74.4%	73.3%	77.4%	5-14	זכר
67.2%	72.9%	76.6%	83.5%	78.9%	71.5%	70.8%	78.8%	71.0%	69.1%	71.2%	77.5%	81.0%	78.3%	70.6%	78.8%	15-24	
70.8%	72.8%	79.4%	84.7%	75.6%	75.5%	71.7%	67.0%	73.2%	64.3%	67.2%	69.9%	81.3%	78.8%	74.2%	78.6%	25-34	
81.1%	71.4%	74.6%	85.2%	72.0%	69.6%	67.7%	67.4%	80.0%	61.2%	72.2%	73.3%	76.6%	73.4%	72.4%	76.1%	35-44	
90.4%	77.1%	83.2%	80.8%	76.6%	73.0%	76.2%	79.2%	77.6%	74.3%	80.9%	76.1%	77.8%	73.7%	70.4%	80.3%	5-14	נקבה
81.0%	80.4%	77.8%	81.5%	79.1%	73.1%	74.9%	75.6%	71.4%	62.7%	68.7%	76.9%	77.1%	73.1%	75.6%	78.0%	15-24	
81.5%	78.1%	83.1%	79.6%	79.6%	70.0%	71.8%	77.1%	73.3%	71.2%	73.7%	76.5%	78.8%	73.6%	73.8%	79.6%	25-34	
79.1%	81.3%	78.0%	87.4%	77.4%	75.0%	71.6%	75.7%	85.5%	75.8%	71.3%	70.9%	74.9%	73.3%	74.6%	79.6%	35-44	

שיעור קבלת חיסון נגד שפעת בקרב חולי אסתמה מתמשכת בני 44-5

28.0%	21.2%	18.9%	25.9%	37.6%	27.6%	20.6%	18.3%	29.3%	45.5%	36.3%	36.1%	35.6%	27.4%	23.6%	33.3%	5-14	זכר
25.0%	14.2%	16.9%	12.8%	22.0%	16.7%	14.3%	10.6%	35.5%	27.7%	29.8%	14.6%	28.4%	18.6%	18.1%	26.8%	15-24	
19.1%	15.4%	15.2%	13.2%	17.4%	15.3%	14.0%	14.2%	25.0%	26.1%	24.8%	20.5%	22.0%	15.5%	19.7%	26.3%	25-34	
21.3%	22.9%	19.3%	26.9%	27.4%	23.3%	18.2%	18.0%	30.0%	33.6%	27.2%	29.7%	29.2%	23.3%	25.3%	35.5%	35-44	
39.8%	28.2%	25.7%	25.3%	31.0%	28.6%	22.4%	21.6%	63.3%	38.1%	39.4%	33.8%	34.3%	26.9%	25.5%	35.2%	5-14	נקבה
17.7%	19.6%	16.7%	19.6%	19.3%	22.1%	23.4%	4.9%	34.3%	40.0%	38.4%	40.0%	28.7%	20.1%	20.6%	36.1%	15-24	
33.7%	29.4%	20.8%	25.5%	27.3%	21.0%	17.3%	13.3%	60.0%	33.3%	34.2%	41.2%	27.0%	23.8%	20.2%	30.9%	25-34	
35.5%	21.4%	20.2%	24.5%	29.1%	22.8%	22.9%	17.1%	35.5%	44.4%	41.9%	43.7%	33.5%	26.8%	27.9%	33.1%	35-44	

שיעור הביצוע של ספירומטריה בקרב חולי COPD ואוכלוסייה בסיכון גבוה ל-COPD בני 74-50

80.3%	73.0%	73.0%	73.7%	52.5%	59.6%	65.2%	57.8%	75.9%	67.7%	73.1%	64.4%	73.8%	72.4%	69.8%	71.9%	50-54	זכר
75.9%	72.4%	77.3%	77.6%	62.6%	69.5%	65.4%	57.4%	79.5%	73.4%	78.0%	71.6%	77.6%	74.3%	73.6%	76.3%	55-59	
85.1%	82.3%	80.8%	78.3%	71.4%	69.3%	68.2%	61.8%	80.4%	77.9%	79.6%	77.6%	79.6%	81.7%	79.8%	79.7%	60-64	
84.0%	82.5%	81.4%	79.1%	68.1%	74.0%	70.1%	62.4%	85.2%	85.6%	82.7%	79.4%	84.1%	82.4%	80.9%	76.8%	65-69	
77.7%	85.3%	82.5%	79.4%	67.8%	77.1%	72.2%	66.5%	85.9%	84.4%	84.6%	81.5%	87.9%	84.4%	83.7%	78.6%	70-74	
68.7%	78.3%	80.5%	66.7%	54.3%	60.5%	60.0%	51.9%	58.9%	59.8%	69.4%	66.4%	70.8%	67.7%	72.5%	70.9%	50-54	נקבה
80.7%	70.8%	76.4%	74.7%	66.8%	64.0%	63.0%	53.0%	76.5%	69.9%	75.1%	79.2%	74.0%	77.7%	76.7%	76.5%	55-59	
78.2%	79.3%	80.2%	71.7%	69.1%	66.9%	67.2%	57.0%	73.3%	75.0%	75.1%	69.0%	79.6%	80.7%	80.6%	75.7%	60-64	
78.3%	82.6%	82.8%	76.5%	71.8%	71.0%	69.5%	60.1%	79.9%	84.0%	79.7%	83.0%	84.1%	82.0%	82.0%	76.4%	65-69	
84.7%	84.9%	79.3%	71.8%	70.7%	75.8%	69.0%	63.8%	79.6%	87.1%	79.5%	83.2%	87.0%	84.9%	81.5%	77.3%	70-74	

תיעוד ערכי בדיקת ספירומטריה בקרב חולי COPD ואוכלוסייה בסיכון גבוה ל-COPD בני 74-50

-	-	-	-	35.7%	29.7%	27.8%	16.5%	58.5%	50.8%	47.3%	44.8%	58.0%	57.4%	60.8%	62.7%	50-54	זכר
-	-	-	-	39.0%	35.7%	32.6%	22.1%	50.0%	52.0%	55.4%	42.0%	53.5%	64.5%	65.3%	65.7%	55-59	
-	-	-	-	41.3%	34.1%	27.5%	16.2%	54.8%	50.2%	54.2%	48.9%	58.1%	67.2%	63.0%	66.9%	60-64	
-	-	-	-	41.5%	32.3%	31.5%	16.3%	56.2%	58.0%	53.0%	42.9%	60.2%	66.9%	66.0%	66.0%	65-69	
-	-	-	-	48.2%	33.8%	30.8%	11.9%	53.6%	50.2%	50.8%	48.8%	56.5%	64.8%	65.8%	67.0%	70-74	
-	-	-	-	42.6%	37.2%	32.9%	13.0%	42.4%	51.3%	43.6%	35.6%	57.8%	57.9%	59.2%	71.6%	50-54	נקבה
-	-	-	-	46.1%	35.8%	29.6%	15.7%	53.8%	48.8%	54.2%	41.5%	56.9%	64.1%	70.2%	63.7%	55-59	
-	-	-	-	43.5%	36.4%	37.2%	18.5%	62.3%	50.2%	51.6%	49.6%	61.4%	62.7%	61.9%	63.6%	60-64	
-	-	-	-	49.2%	37.7%	33.6%	12.8%	60.0%	49.3%	48.0%	52.7%	55.2%	60.1%	67.4%	61.1%	65-69	
-	-	-	-	50.8%	39.6%	37.1%	13.6%	50.5%	45.7%	45.7%	49.4%	57.5%	63.7%	63.7%	58.6%	70-74	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

לב וכלי דם

שיעור הביצוע של בדיקת כולסטרול בקרב בני 35-54

77.5%	75.1%	73.8%	72.0%	79.3%	77.8%	74.5%	73.8%	77.4%	75.0%	73.4%	70.9%	82.6%	79.5%	76.0%	75.3%	35-39	זכר
80.1%	79.2%	78.1%	78.3%	82.2%	81.3%	79.6%	80.2%	80.1%	78.5%	78.7%	77.0%	84.5%	82.8%	80.2%	81.2%	40-44	
84.1%	82.7%	81.7%	81.9%	86.0%	85.0%	82.7%	83.7%	83.1%	82.4%	81.6%	83.2%	87.5%	86.2%	84.5%	84.9%	45-49	
89.2%	88.6%	86.9%	87.2%	90.0%	89.4%	87.7%	87.8%	89.5%	87.5%	86.5%	86.5%	91.2%	90.2%	88.9%	89.5%	50-54	
88.2%	87.1%	85.6%	85.6%	86.7%	86.9%	85.7%	85.7%	88.2%	87.4%	87.1%	81.8%	92.1%	91.3%	90.7%	91.5%	35-39	נקבה
90.6%	90.0%	89.3%	88.8%	88.5%	89.0%	88.1%	88.3%	90.5%	89.8%	89.3%	87.4%	92.5%	92.3%	92.2%	93.0%	40-44	
92.7%	92.7%	91.2%	92.3%	91.6%	91.7%	91.2%	90.8%	92.4%	92.6%	92.2%	91.5%	94.1%	94.0%	93.6%	94.7%	45-49	
94.1%	94.6%	93.5%	94.9%	94.0%	94.3%	93.5%	93.8%	94.1%	94.4%	94.5%	93.7%	95.2%	95.2%	95.3%	96.2%	50-54	

שיעור הביצוע של בדיקת כולסטרול בקרב בני 55-74

61.5%	61.7%	61.0%	63.3%	56.0%	57.7%	57.2%	59.0%	64.4%	60.7%	60.0%	58.9%	64.7%	63.5%	62.5%	63.9%	55-59	זכר
68.8%	68.2%	67.2%	69.5%	64.1%	65.6%	65.0%	65.7%	72.4%	69.2%	67.1%	65.3%	72.5%	71.7%	69.6%	70.8%	60-64	
78.2%	76.8%	74.0%	76.5%	72.3%	73.8%	73.5%	72.6%	78.3%	77.6%	73.8%	72.3%	80.6%	79.2%	77.1%	77.4%	65-69	
82.8%	80.9%	80.0%	80.6%	78.0%	80.0%	78.6%	77.3%	84.4%	81.9%	78.6%	76.4%	85.5%	84.2%	81.4%	81.0%	70-74	
70.6%	69.5%	68.6%	74.3%	56.9%	59.0%	59.6%	63.2%	69.3%	71.2%	71.7%	71.3%	70.5%	71.1%	72.7%	75.6%	55-59	נקבה
73.6%	74.1%	74.1%	77.8%	64.1%	66.5%	68.0%	69.3%	77.3%	76.4%	75.5%	76.2%	76.4%	76.4%	77.5%	80.1%	60-64	
79.0%	78.5%	78.6%	82.3%	71.4%	73.4%	74.0%	73.1%	79.8%	81.5%	80.4%	80.0%	82.4%	82.1%	81.4%	84.3%	65-69	
83.4%	83.0%	81.2%	83.5%	76.9%	78.4%	78.9%	78.7%	85.6%	83.2%	82.5%	83.4%	85.8%	85.4%	84.9%	85.8%	70-74	

שיעור עמידה ביעד LDL לפי קבוצת סיכון למחלות לב בקרב בני 74-35

87.8%	87.5%	87.4%	87.6%	86.2%	84.2%	81.9%	83.9%	83.6%	80.7%	78.5%	79.6%	91.2%	89.6%	88.6%	87.3%	35-44	זכר
82.6%	79.2%	76.7%	76.3%	79.7%	75.8%	73.9%	76.5%	70.0%	65.6%	65.6%	64.6%	81.4%	78.1%	76.9%	75.2%	45-54	
81.5%	79.7%	78.8%	77.8%	82.6%	79.3%	76.6%	76.7%	74.0%	67.7%	65.9%	68.5%	86.2%	82.5%	80.5%	78.4%	55-64	
85.1%	81.1%	79.4%	80.5%	85.8%	82.9%	79.5%	82.1%	75.8%	73.5%	69.7%	70.8%	90.1%	87.4%	85.7%	84.4%	65-74	
96.0%	95.1%	96.5%	96.4%	93.4%	92.2%	92.3%	92.0%	92.0%	91.1%	91.6%	92.7%	96.4%	96.1%	96.1%	96.0%	35-44	נקבה
92.2%	91.2%	91.2%	92.9%	87.5%	85.6%	84.7%	86.6%	81.8%	82.1%	84.0%	85.3%	91.7%	90.4%	90.3%	91.1%	45-54	
83.8%	82.5%	81.1%	84.7%	80.5%	77.6%	75.4%	79.8%	72.4%	71.2%	71.6%	73.4%	85.0%	82.9%	82.6%	84.4%	55-64	
87.2%	85.4%	84.0%	85.8%	84.8%	82.8%	81.0%	83.1%	78.4%	75.9%	74.0%	77.6%	89.2%	88.2%	87.7%	89.2%	65-74	

שיעור בעלי רמת LDL נמוכה או שווה ל-160 מ"ג/ד"ל בקבוצת סיכון נמוך למחלות לב וכלי דם בבני 74-35

91.8%	91.9%	92.4%	93.5%	88.1%	87.0%	85.2%	88.1%	85.8%	83.4%	82.4%	84.1%	92.4%	91.6%	91.6%	90.8%	35-44	זכר
91.7%	90.3%	90.7%	91.7%	86.2%	84.2%	82.5%	88.0%	80.1%	78.9%	78.5%	81.1%	90.0%	88.7%	89.3%	88.6%	45-54	
91.8%	91.9%	92.2%	90.3%	89.1%	87.4%	86.7%	87.8%	82.9%	79.4%	81.3%	84.1%	92.6%	91.8%	91.2%	90.5%	55-64	
95.6%	95.3%	94.1%	94.2%	92.8%	91.4%	87.9%	90.1%	86.5%	86.3%	84.8%	85.6%	96.0%	94.7%	94.6%	94.6%	65-74	
96.7%	95.6%	97.0%	96.9%	93.6%	92.7%	93.0%	92.4%	92.6%	91.8%	92.3%	93.0%	96.6%	96.5%	96.6%	96.3%	35-44	נקבה
93.1%	92.5%	93.1%	94.2%	88.2%	86.8%	86.3%	87.8%	82.6%	83.0%	84.9%	86.1%	92.3%	91.3%	91.6%	92.1%	45-54	
89.5%	89.3%	89.0%	91.7%	84.8%	83.0%	82.6%	85.9%	79.1%	78.9%	79.5%	82.1%	90.1%	89.7%	89.8%	90.5%	55-64	
91.4%	91.1%	89.9%	91.4%	88.6%	87.7%	86.3%	88.3%	83.5%	81.9%	80.8%	85.8%	92.3%	92.2%	92.4%	94.0%	65-74	

שיעור בעלי רמת LDL נמוכה או שווה ל-130 מ"ג/ד"ל בקרב בעלי 2 או יותר נקודות גורמי סיכון למחלות לב וכלי דם ו-Framingham נמוך או שווה ל-20% בבני 74-35

67.5%	70.1%	70.9%	74.8%	71.0%	67.1%	67.8%	67.3%	65.1%	64.3%	63.3%	63.2%	75.3%	73.0%	72.0%	72.5%	35-44	זכר
72.2%	70.8%	69.2%	71.0%	69.9%	68.4%	69.0%	70.3%	60.1%	57.1%	60.4%	58.8%	71.9%	71.2%	72.1%	71.7%	45-54	
75.5%	74.1%	75.0%	77.2%	77.5%	76.2%	75.0%	76.0%	71.9%	65.9%	65.2%	70.0%	82.8%	80.6%	79.9%	79.8%	55-64	
84.4%	80.9%	80.7%	82.1%	85.3%	83.2%	80.9%	82.5%	76.1%	75.3%	72.7%	75.5%	89.1%	87.4%	85.9%	84.2%	65-74	
67.1%	77.1%	80.0%	79.2%	76.4%	75.8%	72.7%	79.4%	71.8%	70.0%	68.3%	83.1%	80.4%	78.5%	80.0%	81.5%	35-44	נקבה
67.7%	69.6%	65.0%	74.2%	65.0%	64.7%	63.0%	70.4%	58.6%	61.3%	65.8%	67.3%	72.0%	71.9%	72.4%	76.9%	45-54	
64.7%	65.8%	66.1%	71.8%	61.6%	62.2%	60.0%	67.1%	53.9%	54.4%	57.3%	59.4%	69.2%	68.8%	70.3%	75.1%	55-64	
77.5%	75.3%	75.6%	78.7%	75.1%	73.6%	72.8%	75.5%	68.6%	67.1%	65.6%	68.0%	81.4%	80.4%	80.5%	83.1%	65-74	

שיעור התייעוד של מדידות לחץ דם בקרב בני 20-54

71.7%	68.4%	64.7%	69.6%	75.5%	76.5%	70.5%	69.1%	57.1%	64.4%	66.9%	69.0%	77.7%	77.7%	77.2%	81.3%	20-24	זכר
71.9%	72.0%	69.5%	73.7%	79.2%	81.1%	79.3%	75.3%	71.3%	74.0%	72.9%	73.5%	75.8%	78.0%	77.9%	80.9%	25-34	
82.5%	80.9%	79.2%	81.5%	85.8%	86.0%	84.7%	82.7%	74.0%	77.3%	77.6%	78.8%	82.0%	83.6%	84.2%	87.6%	35-44	
92.3%	91.3%	89.7%	91.2%	90.9%	91.6%	90.8%	90.1%	82.5%	84.2%	84.4%	87.8%	91.9%	93.0%	94.0%	96.0%	45-54	
83.4%	81.1%	82.8%	87.0%	84.4%	86.5%	85.6%	85.3%	77.1%	81.3%	82.6%	86.0%	86.0%	86.6%	87.1%	89.8%	20-24	נקבה
87.9%	90.0%	92.6%	95.0%	90.7%	92.7%	93.0%	93.2%	87.3%	89.6%	91.2%	92.6%	89.8%	91.4%	91.7%	92.1%	25-34	
92.8%	91.5%	93.0%	94.8%	91.6%	92.3%	92.1%	92.5%	86.6%	87.5%	89.3%	91.4%	90.5%	91.6%	92.4%	94.0%	35-44	
94.5%	94.3%	93.6%	95.7%	91.9%	93.2%	93.7%	93.9%	87.9%	88.5%	89.9%	90.8%	93.9%	95.8%	96.6%	98.0%	45-54	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור התייעוד של מדידות לחץ דם בקרב בני 74-55

59.4%	60.7%	62.7%	67.4%	55.3%	58.3%	59.0%	61.9%	53.8%	53.9%	57.7%	61.8%	59.4%	63.3%	64.9%	69.4%	55-59	זכר
67.1%	68.5%	69.1%	73.3%	62.2%	64.6%	66.1%	68.8%	59.3%	62.1%	63.2%	65.8%	67.6%	71.7%	72.0%	75.1%	60-64	
75.0%	77.1%	75.9%	78.9%	69.7%	72.6%	73.7%	75.6%	68.4%	71.1%	69.9%	72.5%	76.7%	79.2%	78.7%	81.1%	65-69	
83.0%	81.7%	80.6%	82.8%	75.3%	78.9%	79.1%	79.1%	73.2%	74.6%	72.9%	74.8%	82.3%	84.2%	83.3%	84.6%	70-74	
62.3%	64.1%	65.9%	72.9%	54.4%	58.9%	62.4%	65.7%	55.5%	60.1%	63.0%	66.9%	60.6%	65.8%	69.9%	75.5%	55-59	נקבה
69.8%	70.6%	71.9%	76.7%	62.3%	66.5%	68.4%	71.1%	62.2%	66.1%	68.1%	71.7%	68.2%	72.8%	75.3%	80.3%	60-64	
77.7%	77.4%	76.8%	81.3%	68.8%	72.6%	74.1%	76.1%	71.2%	71.9%	73.1%	75.3%	76.3%	79.6%	80.5%	84.8%	65-69	
82.4%	82.4%	81.2%	83.0%	74.7%	77.3%	79.9%	80.0%	74.2%	76.4%	74.7%	77.7%	81.9%	84.2%	84.7%	86.9%	70-74	

שיעור בעלי רמת LDL נמוכה או שווה ל-70 מ"ג/ד"ל או טיפול בסטטינים בעצמות גבוהה לאחר צנתור לבבי / ניתוח מעקפים בקרב בני 79-35

77.5%	82.8%	83.4%	80.9%	78.7%	71.1%	69.1%	62.0%	64.7%	71.1%	57.7%	62.5%	74.7%	73.3%	65.8%	63.8%	35-49	זכר
86.2%	86.7%	85.0%	82.7%	78.4%	74.0%	67.7%	68.2%	76.2%	65.6%	59.6%	61.6%	79.8%	75.1%	70.2%	66.7%	50-59	
88.1%	86.9%	84.1%	83.7%	80.9%	75.8%	74.7%	75.1%	79.8%	75.3%	68.9%	67.5%	84.0%	79.3%	75.1%	72.5%	60-69	
88.6%	86.6%	85.0%	85.4%	81.9%	79.7%	75.2%	78.4%	83.5%	78.1%	69.9%	73.3%	85.8%	83.9%	80.2%	77.0%	70-79	
-	44.7%	33.3%	35.9%	53.7%	58.7%	55.6%	-	-	-	-	-	48.6%	50.0%	53.1%	54.0%	35-49	נקבה
71.8%	62.2%	52.6%	73.4%	75.8%	61.0%	59.7%	58.0%	-	61.7%	46.1%	53.5%	67.3%	69.8%	59.0%	63.5%	50-59	
88.4%	74.1%	79.9%	80.9%	72.5%	67.3%	66.9%	73.9%	69.4%	64.0%	66.2%	68.5%	75.0%	72.9%	72.4%	73.8%	60-69	
79.7%	81.5%	74.6%	75.2%	72.1%	67.2%	67.4%	67.3%	66.0%	68.3%	62.8%	69.8%	75.0%	74.2%	74.6%	74.8%	70-79	

שיעור הטיפול בתרופות להורדת LDL לאחר ניתוח מעקפים ו/או צנתור לבבי בקרב בני 79-35

74.6%	74.6%	78.0%	65.0%	85.4%	77.9%	78.1%	77.3%	86.7%	84.9%	81.1%	72.4%	80.0%	79.4%	74.6%	77.2%	35-49	זכר
80.1%	80.2%	79.3%	76.3%	87.7%	85.2%	79.9%	81.8%	90.1%	82.3%	79.9%	82.3%	85.3%	81.9%	81.3%	79.2%	50-59	
81.4%	82.7%	81.6%	75.1%	90.1%	85.4%	80.9%	85.8%	87.5%	85.7%	83.2%	84.3%	87.3%	85.6%	81.4%	81.4%	60-69	
86.9%	86.0%	86.5%	82.6%	91.3%	88.9%	85.6%	87.3%	90.8%	90.4%	84.6%	85.5%	90.3%	88.9%	86.4%	84.3%	70-79	
-	56.7%	-	-	85.4%	78.6%	69.2%	-	-	-	-	-	60.0%	56.7%	71.4%	78.7%	35-49	נקבה
81.3%	62.7%	57.1%	61.7%	86.9%	83.0%	80.0%	77.5%	-	85.7%	83.3%	-	75.0%	81.1%	78.4%	81.6%	50-59	
78.0%	72.0%	83.9%	77.6%	87.3%	85.9%	87.6%	90.6%	100.0%	86.1%	88.5%	77.3%	86.0%	86.3%	86.1%	87.9%	60-69	
83.9%	81.8%	77.5%	79.5%	90.1%	86.8%	90.0%	86.6%	90.2%	89.3%	88.6%	86.2%	87.7%	88.7%	86.5%	84.2%	70-79	

שיעור ביצוע של בדיקת כולסטרול בדם בקרב מבטחים לאחר צנתור לבבי או ניתוח מעקפים בקרב בני 35-79

-	-	-	-	-	82.6%	75.6%	-	-	-	-	-	60.0%	65.5%	69.3%	63.2%	35-49	זכר
73.9%	80.0%	79.4%	78.1%	85.2%	86.5%	85.8%	86.0%	71.6%	72.2%	72.4%	76.9%	78.7%	78.3%	78.7%	79.3%	50-59	
83.3%	83.8%	84.8%	84.9%	91.5%	91.2%	90.8%	89.9%	82.7%	81.2%	80.8%	82.5%	87.5%	86.3%	84.8%	85.1%	60-69	
90.6%	88.9%	88.3%	87.8%	95.1%	95.5%	93.7%	94.2%	88.3%	87.4%	85.4%	83.3%	91.7%	90.9%	88.9%	89.0%	70-79	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	35-49	נקבה
86.5%	74.7%	82.8%	86.6%	86.8%	90.2%	89.5%	87.3%	-	88.7%	91.8%	84.3%	82.2%	86.8%	87.0%	84.3%	50-59	
91.8%	86.4%	90.4%	91.9%	92.5%	94.8%	93.1%	92.9%	87.5%	86.1%	85.3%	84.0%	88.7%	91.0%	90.3%	92.5%	60-69	
93.7%	90.0%	87.2%	94.0%	94.0%	95.1%	96.4%	96.4%	88.5%	86.1%	89.9%	89.6%	92.8%	93.2%	92.7%	92.3%	70-79	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

סוכרת

שיעור ההימצאות של סוכרת בקרב בני 18 ומעלה

0.9%	0.7%	0.6%	0.4%	0.8%	0.8%	0.6%	0.6%	0.6%	0.4%	0.5%	0.4%	0.9%	0.9%	0.6%	0.5%	18-24	זכר
0.7%	0.6%	0.6%	1.0%	0.5%	0.5%	0.8%	0.9%	0.4%	0.6%	0.7%	0.8%	0.6%	0.7%	0.8%	1.1%	25-34	
1.4%	1.9%	2.6%	4.1%	1.4%	2.0%	2.9%	3.6%	1.8%	2.2%	2.8%	3.6%	1.5%	2.2%	3.3%	4.5%	35-44	
5.0%	6.8%	9.1%	12.6%	4.8%	7.6%	9.8%	12.4%	6.6%	8.9%	9.8%	12.6%	5.5%	8.4%	11.3%	14.3%	45-54	
13.1%	16.5%	19.6%	25.9%	13.1%	17.2%	20.5%	25.2%	18.0%	19.9%	20.8%	24.1%	16.7%	21.0%	23.7%	28.5%	55-64	
25.1%	28.0%	30.2%	37.2%	24.2%	29.4%	32.0%	35.0%	29.1%	31.5%	32.3%	33.0%	29.4%	33.5%	35.5%	40.2%	65-74	
30.7%	32.1%	32.9%	37.1%	30.6%	34.1%	33.5%	36.1%	34.2%	37.7%	35.2%	38.0%	35.1%	38.3%	40.1%	43.2%	75-84	
26.5%	29.9%	26.7%	33.9%	25.9%	28.5%	28.1%	31.1%	28.9%	29.4%	27.8%	26.3%	30.4%	32.0%	32.6%	31.8%	85+	
0.9%	0.6%	0.5%	0.4%	0.8%	0.6%	0.5%	0.5%	0.8%	0.6%	0.5%	0.4%	0.9%	0.7%	0.6%	0.5%	18-24	נקבה
0.7%	0.8%	0.7%	0.8%	0.7%	0.8%	1.0%	1.0%	0.7%	0.7%	0.8%	0.9%	0.6%	0.7%	0.9%	1.1%	25-34	
1.0%	1.4%	1.7%	2.4%	1.1%	1.8%	2.5%	3.0%	1.2%	1.9%	2.1%	2.7%	1.2%	1.6%	2.3%	3.8%	35-44	
2.7%	3.6%	5.3%	9.3%	2.9%	4.6%	7.1%	9.4%	3.8%	5.0%	7.2%	9.6%	3.4%	5.4%	8.3%	13.1%	45-54	
7.5%	10.4%	14.6%	23.5%	7.6%	11.5%	15.7%	21.3%	9.5%	13.1%	17.4%	21.4%	10.4%	14.4%	20.2%	29.6%	55-64	
16.4%	20.1%	26.0%	34.6%	15.4%	22.3%	26.8%	32.1%	19.5%	23.7%	29.3%	34.8%	20.3%	25.4%	32.1%	42.9%	65-74	
22.0%	25.3%	30.4%	37.3%	20.2%	27.2%	31.1%	35.8%	26.1%	30.6%	33.5%	37.0%	26.5%	31.6%	37.2%	44.0%	75-84	
19.5%	23.0%	26.6%	27.7%	18.5%	23.8%	28.3%	30.6%	25.1%	24.7%	28.4%	26.6%	22.9%	26.3%	30.0%	30.9%	85+	

שיעור הביצוע של בדיקת HbA1c בחולי סוכרת בני 18 ומעלה

69.8%	74.2%	78.9%	74.6%	78.9%	78.3%	80.0%	80.6%	78.3%	74.0%	72.6%	70.7%	77.5%	76.5%	72.3%	76.1%	18-34	זכר
73.1%	78.4%	78.6%	83.9%	77.6%	80.6%	81.4%	80.0%	72.3%	75.1%	69.0%	72.5%	76.8%	77.4%	76.0%	78.4%	35-44	
79.3%	82.8%	82.7%	83.9%	83.7%	85.0%	86.0%	86.7%	71.2%	76.8%	76.5%	78.7%	81.2%	82.2%	80.8%	82.4%	45-54	
84.9%	85.7%	86.4%	88.8%	88.9%	89.9%	90.0%	90.3%	85.1%	81.9%	83.6%	83.2%	84.6%	85.4%	85.4%	86.3%	55-64	
91.6%	90.2%	89.2%	91.3%	93.4%	94.5%	94.6%	94.3%	91.3%	89.4%	87.7%	85.4%	90.2%	90.2%	89.8%	90.2%	65-74	
92.6%	91.7%	91.3%	92.2%	95.4%	95.8%	95.4%	95.8%	91.3%	90.7%	89.3%	88.2%	92.1%	91.6%	90.0%	90.5%	75-84	
89.4%	88.7%	86.1%	90.3%	93.7%	93.3%	91.4%	94.2%	90.4%	82.6%	83.7%	81.5%	88.1%	88.5%	88.1%	87.3%	85+	
86.0%	79.9%	83.1%	88.0%	75.7%	74.3%	70.9%	78.5%	77.1%	79.8%	77.8%	77.8%	82.1%	80.3%	80.3%	84.9%	18-34	נקבה
78.4%	84.8%	86.0%	87.9%	70.5%	70.2%	76.0%	83.2%	70.7%	79.5%	82.9%	80.7%	82.8%	80.2%	81.9%	86.9%	35-44	
88.7%	88.1%	87.2%	91.8%	78.0%	79.8%	83.6%	89.2%	83.9%	81.7%	85.9%	87.3%	84.5%	87.0%	85.9%	90.0%	45-54	
87.0%	87.9%	88.6%	92.8%	86.0%	89.6%	91.3%	92.6%	86.0%	86.7%	88.0%	89.6%	88.0%	88.4%	89.1%	91.3%	55-64	
92.1%	92.0%	91.5%	95.3%	94.1%	94.4%	94.3%	94.6%	90.8%	90.5%	90.1%	89.4%	91.1%	91.0%	91.0%	91.9%	65-74	
91.8%	90.5%	91.5%	93.4%	93.9%	94.7%	95.1%	95.5%	91.1%	89.5%	87.9%	86.0%	91.2%	91.2%	91.0%	91.5%	75-84	
88.5%	85.0%	86.2%	89.5%	92.1%	93.1%	92.9%	91.4%	81.2%	85.7%	80.2%	83.2%	84.8%	85.7%	87.0%	88.8%	85+	

שיעור בעלי רמת HbA1c מאוזנת בחולי סוכרת בני 18-84

61.7%	62.7%	52.8%	46.0%	69.5%	62.1%	60.8%	58.0%	83.3%	71.9%	49.6%	48.0%	71.7%	62.5%	51.0%	46.5%	18-34	זכר
67.1%	67.8%	63.4%	49.3%	73.9%	66.4%	60.5%	58.0%	78.7%	62.8%	62.5%	50.2%	72.0%	64.9%	60.0%	51.6%	35-44	
70.2%	63.8%	59.4%	52.6%	70.5%	67.8%	64.0%	62.2%	69.9%	62.3%	61.1%	54.0%	73.7%	67.5%	61.5%	54.3%	45-54	
72.5%	69.8%	65.3%	57.7%	76.1%	72.3%	67.8%	65.3%	77.3%	74.6%	69.1%	57.7%	77.7%	71.9%	67.1%	59.5%	55-64	
81.4%	77.3%	74.9%	66.8%	83.4%	80.2%	77.1%	77.4%	86.4%	81.0%	78.2%	69.2%	86.3%	81.4%	75.6%	69.0%	65-74	
88.6%	86.1%	82.7%	80.8%	91.6%	89.7%	88.8%	87.9%	90.6%	88.4%	88.8%	82.4%	91.7%	89.1%	85.5%	82.2%	75-84	
60.8%	76.2%	72.3%	63.4%	70.7%	73.4%	71.5%	67.9%	77.8%	70.7%	70.0%	64.2%	76.1%	70.1%	63.4%	58.1%	18-34	
78.3%	71.1%	69.3%	60.3%	81.0%	77.7%	76.7%	67.5%	75.9%	71.9%	72.6%	55.7%	79.5%	76.7%	70.3%	62.7%	35-44	
69.7%	73.1%	67.9%	60.8%	80.3%	77.0%	73.1%	71.0%	80.0%	78.7%	71.1%	57.8%	79.4%	74.5%	68.9%	62.6%	45-54	
78.9%	77.3%	72.3%	66.5%	81.8%	80.1%	76.5%	73.3%	81.7%	79.0%	76.1%	66.2%	83.3%	79.1%	74.0%	68.6%	55-64	
84.7%	82.8%	80.4%	73.6%	87.8%	85.8%	82.9%	81.5%	87.6%	85.5%	82.7%	74.7%	89.0%	85.2%	80.2%	74.1%	65-74	
87.7%	84.5%	84.8%	85.2%	93.3%	91.3%	91.3%	86.8%	92.1%	90.5%	89.5%	84.9%	92.2%	90.4%	87.1%	83.1%	75-84	

שיעור בעלי רמת HbA1c גבוהה מ-9% בחולי סוכרת בני 18 ומעלה

5.0%	9.3%	27.2%	26.4%	8.0%	14.5%	15.4%	21.9%	5.6%	14.0%	20.5%	26.0%	7.7%	16.3%	21.7%	28.0%	18-34	זכר
9.2%	9.3%	11.8%	20.2%	9.5%	10.2%	12.8%	18.3%	8.5%	13.8%	14.6%	24.9%	7.9%	13.5%	15.8%	22.4%	35-44	
9.4%	12.7%	14.5%	16.4%	8.0%	10.2%	11.7%	14.4%	12.1%	14.2%	13.3%	19.2%	8.0%	10.3%	15.3%	19.5%	45-54	
7.9%	9.6%	10.2%	12.2%	5.6%	7.7%	9.8%	10.9%	7.4%	7.7%	10.2%	14.5%	7.0%	9.2%	11.4%	15.3%	55-64	
3.6%	4.9%	6.1%	9.7%	3.7%	4.4%	6.4%	5.3%	2.9%	4.9%	6.7%	10.5%	3.3%	4.9%	7.2%	9.8%	65-74	
2.6%	2.9%	5.1%	5.7%	2.8%	3.3%	3.9%	5.0%	2.6%	3.8%	4.2%	9.2%	2.8%	3.7%	5.7%	7.2%	75-84	
1.6%	5.0%	4.1%	10.7%	2.0%	2.9%	4.1%	6.2%	1.8%	2.0%	3.1%	5.2%	3.3%	4.1%	5.1%	7.0%	85+	
8.1%	9.8%	12.9%	14.4%	9.2%	8.4%	9.3%	14.6%	11.1%	6.7%	17.7%	17.9%	6.5%	9.9%	16.3%	20.4%	18-34	נקבה
5.8%	7.0%	12.9%	15.1%	6.0%	8.8%	9.1%	12.3%	6.9%	8.6%	9.4%	17.7%	6.0%	6.7%	12.4%	16.2%	35-44	
9.5%	7.9%	10.3%	12.6%	6.0%	6.4%	9.6%	9.7%	8.0%	6.1%	10.8%	18.4%	6.0%	8.9%	12.4%	15.5%	45-54	
5.6%	6.0%	7.7%	10.3%	4.7%	5.4%	7.3%	9.1%	4.8%	6.6%	8.1%	11.7%	5.3%	6.5%	9.7%	12.1%	55-64	
2.7%	3.9%	5.0%	7.5%	2.5%	3.0%	4.3%	5.2%	2.7%	4.4%	5.1%	8.5%	2.4%	4.4%	6.3%	9.1%	65-74	
3.4%	4.1%	3.7%	5.3%	2.0%	3.3%	3.3%	4.8%	2.8%	3.6%	4.0%	6.0%	2.5%	3.5%	5.4%	7.3%	75-84	
2.6%	2.3%	3.2%	4.3%	3.7%	2.5%	2.6%	4.5%	1.3%	3.7%	2.8%	5.0%	3.8%	4.3%	4.6%	6.0%	85+	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור הביצוע של בדיקות עיניים בחולי סוכרת בני 18-84

46.5%	50.9%	55.1%	54.0%	73.6%	65.6%	69.9%	68.3%	-	36.4%	41.7%	48.3%	58.7%	58.0%	62.6%	62.8%	18-34	זכר
63.5%	56.1%	58.6%	59.2%	72.2%	71.5%	70.0%	68.8%	50.8%	40.9%	37.0%	45.0%	58.6%	62.3%	62.7%	65.4%	35-44	
60.9%	58.9%	59.6%	62.5%	75.0%	76.1%	76.8%	77.2%	41.6%	42.8%	47.6%	51.2%	63.0%	64.2%	67.5%	69.3%	45-54	
65.0%	59.8%	64.5%	66.4%	79.2%	80.1%	78.3%	77.0%	47.7%	48.7%	51.7%	53.5%	65.8%	67.1%	70.3%	71.4%	55-64	
70.1%	70.7%	70.0%	69.6%	84.7%	85.3%	83.4%	82.5%	59.2%	58.4%	58.9%	63.2%	74.0%	75.0%	74.7%	74.5%	65-74	
70.4%	70.9%	71.5%	66.1%	86.2%	83.4%	82.3%	82.1%	63.4%	58.8%	59.5%	58.9%	75.4%	74.6%	72.6%	71.1%	75-84	
55.8%	48.0%	55.1%	62.7%	62.6%	63.9%	57.4%	61.9%	37.1%	41.5%	41.3%	54.4%	62.7%	62.6%	66.9%	69.7%	18-34	נקבה
60.2%	58.2%	63.5%	66.5%	61.3%	65.4%	65.9%	72.4%	41.5%	48.4%	43.9%	48.7%	60.6%	64.6%	67.5%	71.6%	35-44	
69.9%	63.6%	68.1%	72.1%	71.6%	73.4%	74.6%	76.5%	49.7%	46.8%	56.7%	62.1%	68.0%	69.8%	72.6%	75.4%	45-54	
72.9%	66.4%	72.6%	74.0%	79.3%	81.0%	79.5%	77.9%	57.9%	57.5%	59.0%	63.2%	74.3%	73.2%	74.9%	75.8%	55-64	
73.9%	73.6%	70.5%	73.5%	86.1%	84.4%	85.4%	81.6%	64.5%	66.1%	65.9%	64.2%	79.4%	77.7%	76.0%	75.4%	65-74	
71.0%	65.6%	66.6%	60.9%	83.6%	80.6%	78.0%	77.5%	61.1%	60.3%	56.8%	56.9%	76.2%	73.9%	70.1%	66.5%	75-84	

שיעור הביצוע של בדיקות לאיתור חלבון בשתן בחולי סוכרת בני 18-84

66.3%	61.0%	69.6%	64.8%	70.3%	63.4%	69.4%	68.3%	-	63.6%	54.3%	57.5%	61.8%	60.7%	60.1%	65.1%	18-34	זכר
67.3%	70.8%	67.9%	74.6%	69.7%	73.2%	70.4%	64.3%	63.1%	61.7%	55.3%	59.2%	63.6%	64.4%	65.1%	66.9%	35-44	
73.5%	76.4%	74.7%	74.4%	76.1%	75.7%	75.9%	75.0%	64.2%	66.3%	62.9%	65.5%	67.7%	70.6%	70.2%	71.1%	45-54	
75.9%	76.3%	77.4%	79.8%	81.3%	80.9%	79.6%	80.1%	76.6%	73.0%	70.5%	70.1%	73.4%	74.6%	75.6%	76.5%	55-64	
82.0%	80.7%	78.8%	82.5%	85.9%	85.3%	85.0%	82.6%	83.5%	78.3%	74.6%	73.3%	80.3%	80.4%	80.7%	80.1%	65-74	
79.9%	79.9%	79.5%	75.9%	86.2%	85.1%	83.1%	82.9%	82.5%	77.0%	73.3%	69.8%	80.9%	81.1%	79.1%	79.2%	75-84	
72.1%	56.4%	66.7%	70.2%	62.6%	60.5%	55.7%	57.4%	57.1%	60.6%	50.9%	51.9%	58.6%	58.3%	56.1%	64.7%	18-34	נקבה
73.9%	70.9%	68.7%	73.3%	58.1%	56.9%	62.9%	67.6%	51.2%	63.4%	58.0%	58.8%	60.3%	60.3%	63.3%	71.1%	35-44	
73.8%	76.8%	73.3%	78.2%	66.8%	68.6%	72.5%	77.8%	71.1%	63.6%	66.9%	70.0%	63.6%	70.6%	71.6%	76.8%	45-54	
74.7%	76.3%	78.0%	82.8%	77.5%	80.1%	82.0%	84.2%	73.0%	74.6%	71.3%	72.2%	72.1%	74.4%	77.2%	81.1%	55-64	
79.6%	80.6%	79.7%	82.3%	86.2%	85.3%	85.0%	84.8%	78.4%	77.0%	74.5%	69.9%	77.6%	77.9%	79.2%	80.7%	65-74	
72.4%	73.4%	73.2%	74.8%	83.3%	82.5%	81.1%	80.5%	71.9%	71.2%	67.4%	65.8%	74.8%	76.5%	75.1%	74.4%	75-84	

שיעור התיעדוד של GFR בחולי סוכרת בני 18-84

72.1%	74.2%	78.5%	75.8%	81.7%	79.0%	81.7%	80.9%	-	71.4%	70.3%	70.7%	80.1%	79.4%	75.3%	79.0%	18-34	זכר
74.0%	80.7%	79.1%	84.3%	81.9%	82.5%	82.5%	80.6%	72.3%	74.1%	70.4%	71.6%	80.2%	79.8%	78.1%	80.7%	35-44	
80.7%	83.3%	83.4%	84.1%	84.9%	86.3%	86.8%	87.2%	69.5%	77.3%	76.9%	79.0%	83.1%	84.3%	83.1%	84.3%	45-54	
86.4%	87.4%	87.7%	89.2%	90.0%	91.2%	91.2%	91.5%	84.9%	81.8%	83.9%	83.1%	86.3%	87.6%	87.8%	88.2%	55-64	
93.3%	92.5%	91.0%	92.3%	94.7%	95.6%	95.5%	95.3%	92.5%	90.1%	88.4%	86.6%	92.9%	92.8%	92.2%	92.2%	65-74	
95.0%	94.6%	94.1%	93.5%	96.8%	97.1%	97.1%	97.6%	91.4%	92.7%	91.2%	90.4%	95.8%	95.0%	93.7%	93.4%	75-84	
89.5%	82.1%	86.0%	87.3%	85.2%	83.4%	81.9%	87.0%	74.3%	83.0%	76.0%	78.5%	85.0%	83.3%	84.9%	88.8%	18-34	נקבה
83.0%	86.1%	87.0%	87.9%	80.1%	81.3%	84.6%	89.5%	68.3%	82.6%	82.5%	79.0%	87.1%	83.8%	84.7%	89.6%	35-44	
89.3%	89.6%	88.3%	92.0%	85.3%	86.0%	88.1%	91.9%	82.6%	81.0%	87.0%	87.3%	86.7%	89.9%	88.5%	91.4%	45-54	
88.2%	89.7%	89.9%	93.4%	89.1%	92.1%	92.8%	93.8%	87.1%	87.2%	88.0%	90.1%	90.4%	91.0%	91.3%	92.8%	55-64	
93.2%	93.5%	92.9%	95.9%	95.6%	95.7%	95.7%	95.3%	91.2%	91.5%	90.7%	89.6%	93.7%	93.5%	93.5%	94.2%	65-74	
93.5%	93.6%	93.8%	94.2%	96.1%	96.6%	96.5%	96.9%	93.0%	90.9%	90.3%	89.5%	95.4%	95.0%	94.5%	94.4%	75-84	

שיעור חולי סוכרת עם פגיעה כלייתית בני 18-84

2.9%	12.0%	13.1%	16.8%	5.5%	6.7%	8.2%	12.3%	-	12.1%	16.3%	15.5%	7.5%	11.9%	14.4%	18.9%	18-34	זכר
12.2%	19.7%	23.9%	24.5%	15.3%	15.6%	15.9%	18.9%	22.8%	25.9%	23.6%	23.6%	14.2%	18.5%	22.4%	24.6%	35-44	
20.6%	25.0%	27.2%	27.6%	15.4%	17.7%	21.5%	22.9%	28.4%	25.0%	24.0%	26.1%	16.4%	21.0%	25.6%	26.9%	45-54	
28.2%	32.0%	31.7%	34.8%	19.5%	22.3%	25.1%	26.5%	28.7%	25.5%	28.7%	32.5%	23.8%	28.2%	31.1%	32.9%	55-64	
38.3%	42.3%	42.4%	44.4%	33.5%	33.7%	36.2%	35.1%	33.9%	31.6%	33.2%	36.8%	34.9%	37.5%	40.0%	42.5%	65-74	
57.0%	58.6%	59.9%	58.2%	51.3%	51.6%	53.2%	49.9%	40.2%	41.5%	39.9%	46.0%	50.8%	52.7%	54.4%	55.4%	75-84	
5.3%	12.5%	15.5%	15.5%	5.0%	5.6%	7.2%	11.3%	9.7%	14.6%	11.5%	17.1%	8.7%	13.4%	13.0%	20.9%	18-34	נקבה
12.2%	18.4%	17.2%	21.3%	5.9%	8.6%	11.9%	11.8%	9.4%	13.2%	15.9%	20.3%	11.1%	15.5%	17.9%	20.8%	35-44	
16.4%	19.5%	22.6%	22.8%	11.0%	12.6%	14.6%	16.1%	17.6%	13.6%	18.1%	24.8%	13.1%	16.7%	20.5%	22.9%	45-54	
16.4%	21.0%	23.8%	27.8%	13.7%	15.4%	17.0%	19.1%	16.0%	16.3%	19.1%	24.8%	15.8%	19.5%	23.4%	24.9%	55-64	
30.9%	32.1%	34.5%	37.9%	23.1%	25.3%	27.5%	27.1%	20.2%	21.1%	22.2%	23.7%	25.2%	28.2%	32.3%	35.5%	65-74	
52.7%	53.3%	54.8%	56.8%	41.9%	44.4%	47.9%	49.7%	27.6%	26.6%	27.0%	30.5%	43.7%	46.8%	51.0%	53.2%	75-84	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור התיעוד של מדידות לחץ דם בחולי סוכרת בני 18 ומעלה

61.6%	62.3%	64.8%	72.5%	69.5%	63.2%	67.2%	70.5%	-	37.7%	58.3%	64.4%	61.0%	61.6%	67.3%	73.9%	18-34
72.1%	72.8%	75.9%	79.2%	65.7%	66.3%	71.7%	74.1%	53.8%	57.5%	51.1%	63.6%	63.8%	69.1%	72.4%	77.5%	35-44
76.0%	75.8%	80.8%	83.2%	69.4%	71.1%	75.0%	78.6%	54.7%	62.9%	66.9%	74.9%	71.4%	76.7%	78.3%	82.3%	45-54
83.0%	81.7%	85.1%	88.9%	75.0%	77.8%	79.4%	82.6%	71.1%	74.1%	75.8%	79.9%	80.6%	84.3%	85.1%	88.5%	55-64
88.1%	88.7%	88.1%	91.5%	82.1%	84.9%	86.3%	88.5%	81.8%	80.1%	80.2%	84.6%	88.1%	89.6%	89.6%	91.3%	65-74
90.7%	90.6%	90.8%	90.0%	86.6%	88.9%	89.0%	90.4%	81.2%	79.2%	76.0%	82.7%	90.8%	92.5%	91.6%	92.5%	75-84
87.0%	89.7%	82.9%	89.0%	87.1%	87.8%	87.6%	83.9%	79.2%	76.5%	69.0%	69.7%	89.9%	91.2%	91.3%	92.1%	85+
72.1%	69.3%	79.4%	79.5%	70.6%	68.6%	68.9%	70.7%	68.6%	55.3%	55.1%	66.5%	62.0%	64.6%	70.9%	80.4%	18-34
73.9%	76.8%	81.1%	83.7%	62.3%	63.8%	70.4%	75.9%	51.2%	64.6%	65.8%	68.5%	65.8%	70.2%	72.6%	83.7%	35-44
72.8%	79.8%	82.2%	86.4%	67.2%	70.8%	75.7%	80.3%	69.1%	63.6%	73.3%	79.6%	73.4%	79.2%	82.6%	87.5%	45-54
82.2%	83.9%	85.7%	90.8%	73.6%	78.7%	81.2%	82.5%	70.8%	77.0%	79.2%	83.6%	81.6%	85.1%	87.3%	90.8%	55-64
90.0%	89.2%	89.8%	92.5%	82.3%	84.7%	86.4%	87.6%	81.3%	81.3%	81.7%	83.3%	88.0%	89.2%	90.4%	92.3%	65-74
89.6%	88.5%	88.5%	89.6%	84.7%	86.0%	87.9%	91.7%	77.2%	79.6%	75.9%	78.0%	89.8%	91.1%	91.3%	92.3%	75-84
80.2%	81.7%	80.1%	82.9%	82.9%	82.3%	84.7%	86.6%	68.0%	69.0%	66.9%	66.5%	84.3%	87.1%	87.7%	87.5%	85+

שיעור בעלי לחץ דם נמוך או שווה ל-140/90 מ"מ"כ בחולי סוכרת בני 18 ומעלה

94.3%	91.9%	95.6%	90.9%	94.5%	91.2%	93.2%	93.9%	-	-	93.1%	88.4%	93.6%	93.0%	94.5%	94.9%	18-34
88.0%	85.8%	81.6%	84.3%	81.7%	81.9%	83.9%	87.6%	85.7%	80.2%	86.0%	88.8%	85.0%	86.5%	84.7%	89.0%	35-44
78.9%	80.8%	79.0%	83.7%	80.2%	80.1%	81.0%	84.2%	86.5%	79.2%	79.1%	84.8%	82.4%	82.7%	82.1%	84.7%	45-54
78.1%	77.0%	77.1%	78.9%	79.6%	79.0%	78.6%	81.3%	78.1%	79.8%	78.2%	83.6%	80.0%	77.2%	77.3%	81.2%	55-64
76.8%	74.5%	73.8%	75.6%	78.3%	77.5%	78.4%	79.5%	80.4%	79.9%	78.5%	79.8%	77.9%	76.4%	75.3%	78.3%	65-74
74.1%	75.0%	74.7%	74.7%	75.5%	76.8%	78.4%	77.2%	82.2%	79.6%	81.6%	85.4%	77.3%	75.0%	75.1%	78.2%	75-84
70.7%	73.3%	78.5%	78.3%	72.1%	74.3%	77.5%	78.6%	82.8%	79.4%	82.9%	75.9%	75.7%	75.4%	74.0%	77.8%	85+
93.5%	96.0%	97.9%	95.7%	95.9%	95.8%	94.8%	95.3%	-	98.1%	97.8%	98.1%	96.5%	94.2%	95.8%	96.1%	18-34
90.8%	89.6%	88.4%	89.7%	90.5%	89.1%	87.5%	91.8%	-	95.2%	89.8%	93.3%	90.2%	89.8%	90.6%	91.1%	35-44
84.9%	84.4%	82.2%	85.5%	85.5%	84.6%	85.8%	87.9%	90.3%	82.9%	88.3%	85.2%	85.5%	84.6%	84.1%	86.4%	45-54
84.4%	82.0%	81.9%	81.4%	81.9%	83.1%	82.0%	83.7%	83.7%	80.2%	82.8%	84.4%	84.2%	82.8%	81.7%	83.0%	55-64
76.1%	75.8%	75.6%	76.3%	76.9%	78.8%	78.7%	80.1%	79.3%	80.1%	80.4%	83.1%	79.3%	78.4%	77.5%	78.0%	65-74
72.3%	69.6%	68.6%	73.2%	73.5%	73.1%	74.3%	73.3%	79.1%	80.3%	78.1%	74.9%	74.2%	72.5%	72.4%	75.3%	75-84
66.1%	66.2%	68.6%	69.0%	65.3%	66.5%	69.4%	74.3%	78.4%	77.2%	73.7%	79.5%	71.9%	71.6%	71.5%	74.0%	85+

שיעור הביצוע של בדיקת כולסטרול בחולי סוכרת בני 18 ומעלה

72.1%	73.0%	78.1%	74.9%	77.7%	75.7%	79.3%	78.4%	-	74.0%	70.3%	70.7%	76.0%	76.1%	71.8%	75.0%	18-34
74.0%	79.7%	79.3%	83.5%	78.9%	80.3%	80.9%	78.2%	70.8%	74.1%	69.3%	69.8%	78.5%	78.4%	76.5%	79.3%	35-44
80.4%	83.2%	83.0%	84.0%	83.1%	84.6%	84.8%	86.3%	70.0%	74.7%	74.4%	78.2%	81.6%	82.9%	81.6%	82.9%	45-54
85.9%	87.0%	87.1%	88.9%	87.9%	89.4%	89.2%	90.0%	83.7%	81.2%	83.1%	81.9%	85.2%	86.1%	86.2%	86.9%	55-64
92.0%	91.3%	90.3%	91.7%	92.8%	94.2%	94.0%	93.8%	91.2%	89.0%	87.3%	85.5%	91.5%	91.4%	90.9%	90.9%	65-74
94.1%	92.9%	92.8%	91.9%	95.0%	95.2%	95.2%	94.6%	90.3%	89.7%	87.9%	87.7%	94.1%	93.4%	92.0%	91.7%	75-84
92.3%	90.8%	89.0%	91.7%	92.6%	91.3%	90.6%	92.4%	84.8%	83.2%	82.4%	75.6%	91.7%	91.2%	90.6%	89.2%	85+
88.4%	78.2%	84.0%	87.0%	77.4%	72.8%	68.3%	73.0%	71.4%	76.6%	70.7%	72.8%	80.6%	77.4%	79.8%	83.3%	18-34
83.0%	84.4%	85.3%	87.2%	69.3%	72.6%	74.3%	84.3%	70.7%	77.6%	79.2%	77.3%	82.6%	80.1%	83.0%	87.7%	35-44
89.3%	89.1%	87.9%	92.0%	80.4%	81.5%	83.1%	88.4%	83.2%	81.7%	86.1%	87.6%	85.3%	88.1%	87.1%	90.7%	45-54
87.0%	89.2%	89.9%	92.9%	86.5%	90.0%	90.6%	92.0%	85.7%	86.7%	88.0%	89.0%	89.0%	89.7%	90.5%	91.9%	55-64
92.5%	92.7%	92.2%	95.3%	93.6%	94.1%	94.0%	93.1%	90.4%	90.4%	89.9%	89.0%	92.5%	92.3%	92.4%	92.8%	65-74
92.2%	92.4%	92.6%	93.4%	93.1%	94.2%	93.8%	94.6%	90.4%	88.5%	88.3%	85.2%	93.3%	93.1%	92.6%	92.5%	75-84
90.8%	85.9%	87.4%	90.0%	86.7%	89.3%	89.8%	89.0%	78.7%	82.8%	78.9%	82.7%	88.5%	89.2%	89.4%	90.4%	85+

שיעור בעלי רמת LDL נמוכה או שווה ל-100 מ"ג/ד"ל בחולי סוכרת בני 18-79

79.1%	83.1%	75.5%	69.7%	58.7%	61.7%	57.9%	64.5%	-	-	47.9%	46.8%	61.9%	62.4%	63.0%	56.4%	18-29
69.2%	62.3%	61.3%	57.0%	42.9%	45.7%	52.6%	51.6%	-	42.0%	37.5%	42.6%	47.8%	50.8%	53.8%	55.6%	30-39
69.0%	60.0%	60.6%	58.0%	56.1%	55.6%	55.7%	56.2%	51.7%	45.9%	42.3%	41.7%	57.2%	57.6%	58.1%	58.6%	40-49
69.9%	69.3%	69.2%	66.3%	69.0%	67.6%	65.1%	66.8%	52.4%	56.0%	54.2%	51.9%	69.2%	68.6%	67.4%	67.6%	50-59
82.5%	78.2%	76.2%	77.2%	79.9%	76.1%	75.5%	76.6%	67.9%	66.1%	62.7%	60.7%	80.3%	78.5%	75.7%	75.4%	60-69
87.8%	86.0%	85.5%	84.9%	86.2%	84.3%	82.1%	83.3%	77.0%	71.5%	70.1%	68.8%	86.0%	83.9%	82.1%	80.2%	70-79
70.4%	73.5%	71.6%	70.5%	49.8%	55.2%	56.0%	46.7%	-	52.4%	41.9%	52.2%	54.1%	53.8%	57.8%	56.9%	18-29
54.7%	54.8%	58.3%	57.3%	49.3%	43.0%	42.5%	44.2%	-	28.9%	27.0%	26.3%	50.6%	52.5%	48.7%	52.5%	30-39
53.3%	52.2%	55.8%	60.1%	47.5%	44.3%	46.1%	51.5%	40.3%	33.3%	36.2%	39.5%	50.4%	51.8%	55.3%	57.6%	40-49
56.8%	56.4%	60.5%	63.1%	51.2%	51.2%	52.7%	60.2%	37.5%	42.4%	45.1%	45.9%	56.6%	58.1%	59.5%	64.6%	50-59
69.4%	69.2%	70.8%	73.8%	65.6%	65.2%	65.9%	68.6%	56.2%	53.7%	54.3%	55.9%	68.5%	69.9%	70.2%	72.7%	60-69
77.4%	76.6%	76.3%	78.7%	74.0%	73.9%	71.2%	75.3%	61.0%	61.8%	59.5%	60.9%	74.7%	74.9%	74.7%	76.5%	70-79

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור התיעוד של מרכיבי BMI בחולי סוכרת בני 20-64

69.3%	67.1%	69.7%	75.6%	67.2%	61.0%	66.5%	71.3%	-	48.6%	56.5%	66.9%	63.9%	65.7%	69.9%	75.5%	20-34	זכר
76.0%	78.1%	77.8%	81.8%	61.9%	65.7%	66.2%	70.4%	60.0%	64.8%	53.2%	66.6%	71.3%	72.8%	75.1%	76.5%	35-44	
80.9%	80.6%	82.8%	82.7%	64.0%	66.5%	69.0%	73.5%	61.3%	66.0%	69.3%	74.4%	74.5%	78.0%	77.7%	80.6%	45-54	
85.6%	82.0%	83.5%	86.7%	65.6%	68.3%	71.5%	74.7%	69.7%	72.4%	72.8%	76.6%	81.5%	84.8%	84.6%	87.4%	55-64	
74.0%	74.4%	83.8%	87.1%	69.2%	69.7%	65.5%	65.9%	66.7%	60.7%	66.4%	70.7%	67.8%	72.9%	76.2%	81.4%	20-34	נקבה
83.0%	77.2%	84.0%	87.4%	61.0%	64.2%	69.3%	75.7%	58.5%	67.7%	70.6%	73.1%	70.3%	75.8%	76.6%	83.6%	35-44	
81.9%	83.3%	83.1%	87.8%	63.7%	66.9%	71.0%	76.3%	75.8%	72.3%	77.3%	80.5%	77.2%	81.5%	83.4%	85.9%	45-54	
82.2%	84.6%	86.5%	89.6%	64.9%	69.8%	72.8%	75.9%	76.0%	73.7%	78.0%	80.3%	83.1%	85.9%	87.3%	89.6%	55-64	

שיעור ההימצאות של עודף משקל בקרב חולי סוכרת בני 20-64

32.7%	28.6%	22.4%	26.3%	30.3%	26.1%	29.8%	21.0%	-	29.4%	23.0%	22.4%	30.7%	29.7%	32.3%	28.9%	20-34	זכר
34.2%	37.4%	31.0%	30.2%	33.0%	36.5%	32.2%	28.5%	33.3%	32.8%	36.3%	34.7%	39.1%	33.1%	34.0%	34.8%	35-44	
41.1%	36.8%	36.3%	35.7%	39.3%	37.8%	34.8%	34.1%	40.9%	39.2%	37.2%	36.3%	38.3%	38.7%	38.1%	38.9%	45-54	
47.4%	41.9%	40.2%	41.4%	46.9%	41.5%	39.8%	39.5%	45.9%	43.8%	40.7%	43.3%	46.0%	43.4%	42.5%	42.4%	55-64	
22.2%	30.3%	21.5%	27.8%	28.1%	26.8%	31.8%	27.7%	-	25.9%	33.7%	29.8%	24.7%	30.5%	24.5%	25.1%	20-34	נקבה
37.0%	25.1%	29.1%	28.2%	25.6%	27.6%	25.5%	27.1%	-	24.8%	30.5%	18.4%	25.7%	26.4%	25.9%	25.6%	35-44	
34.8%	32.1%	27.4%	24.3%	31.8%	30.9%	29.7%	24.8%	30.1%	25.3%	24.8%	21.7%	31.2%	31.8%	29.5%	25.4%	45-54	
35.5%	33.7%	30.7%	25.1%	35.9%	32.7%	30.8%	27.7%	30.4%	35.9%	30.7%	28.9%	37.4%	35.0%	31.5%	27.0%	55-64	

שיעור ההימצאות של השמנת יתר בקרב חולי סוכרת בני 20-64

11.5%	23.5%	17.8%	20.8%	12.0%	16.9%	22.6%	22.2%	-	20.6%	16.1%	20.6%	13.8%	19.1%	17.0%	22.3%	20-34	זכר
21.5%	26.8%	28.0%	32.4%	26.5%	27.9%	29.6%	35.0%	33.3%	30.4%	30.3%	30.2%	25.1%	29.6%	28.2%	31.5%	35-44	
30.1%	32.4%	32.2%	33.4%	31.6%	32.7%	35.0%	33.7%	28.2%	31.1%	31.7%	33.1%	31.0%	29.7%	30.1%	32.0%	45-54	
27.2%	31.1%	31.9%	30.6%	27.0%	30.1%	31.9%	32.7%	24.8%	26.5%	30.8%	29.4%	26.9%	28.3%	27.5%	28.6%	55-64	
14.8%	14.3%	28.2%	20.9%	15.7%	21.9%	24.6%	25.9%	-	14.8%	25.7%	23.1%	17.1%	16.5%	20.1%	24.0%	20-34	נקבה
23.3%	29.0%	29.8%	32.2%	24.5%	27.6%	27.8%	25.4%	-	27.5%	26.3%	28.7%	24.1%	27.4%	29.0%	30.7%	35-44	
27.3%	29.6%	29.4%	35.5%	27.6%	29.4%	31.3%	34.4%	35.4%	32.5%	34.1%	35.5%	29.2%	27.0%	30.3%	34.0%	45-54	
27.9%	30.2%	30.7%	34.6%	27.7%	30.5%	30.7%	33.6%	35.1%	28.5%	31.9%	33.4%	27.1%	29.2%	30.9%	33.3%	55-64	

שיעור ההימצאות של השמנת יתר חולנית בקרב חולי סוכרת בני 20-64

13.5%	11.2%	14.5%	22.9%	14.5%	15.3%	18.3%	16.5%	-	23.5%	26.4%	26.2%	14.2%	15.3%	14.9%	17.4%	20-34	זכר
21.5%	23.0%	26.1%	26.9%	20.0%	22.5%	23.9%	25.1%	15.4%	24.0%	21.9%	25.8%	17.2%	19.7%	20.9%	20.2%	35-44	
16.0%	17.6%	21.4%	20.6%	16.4%	19.3%	19.6%	23.1%	14.1%	17.4%	19.2%	19.9%	15.3%	17.0%	17.2%	16.1%	45-54	
10.0%	13.1%	15.3%	13.9%	9.9%	13.9%	15.2%	15.6%	8.8%	12.4%	13.5%	13.7%	9.5%	10.4%	11.4%	12.2%	55-64	
18.5%	22.7%	21.5%	23.0%	15.1%	18.2%	24.6%	25.9%	-	31.5%	18.8%	25.0%	18.7%	22.2%	25.7%	25.2%	20-34	נקבה
32.9%	28.4%	31.0%	29.8%	24.0%	26.1%	32.5%	35.7%	-	33.0%	31.1%	44.8%	23.4%	26.6%	30.5%	33.9%	35-44	
23.3%	26.7%	34.1%	31.7%	22.0%	24.8%	27.7%	34.1%	19.5%	29.2%	31.0%	35.7%	22.9%	25.9%	27.9%	33.0%	45-54	
17.2%	22.9%	28.7%	32.7%	17.0%	22.0%	27.0%	29.8%	15.9%	21.4%	27.0%	28.7%	16.8%	19.6%	24.6%	31.2%	55-64	

שיעור התיעוד של מרכיבי BMI בחולי סוכרת בני 65-84

84.7%	85.9%	83.9%	88.4%	68.8%	72.1%	73.4%	78.1%	77.3%	73.5%	77.9%	77.4%	83.8%	84.9%	84.5%	87.9%	65-69	זכר
87.8%	88.4%	86.3%	87.1%	70.9%	75.5%	76.6%	78.8%	87.6%	82.3%	78.7%	83.0%	88.3%	88.8%	88.8%	89.5%	70-74	
87.1%	88.1%	89.6%	85.2%	72.9%	76.9%	78.1%	79.5%	76.7%	75.8%	70.8%	70.5%	88.7%	89.9%	87.4%	89.1%	75-79	
85.9%	81.7%	83.6%	77.0%	74.1%	73.7%	76.3%	75.1%	73.4%	68.6%	65.0%	69.8%	89.1%	88.4%	87.5%	87.0%	80-84	
87.5%	84.7%	88.8%	90.2%	70.2%	73.4%	75.3%	78.8%	81.5%	75.8%	76.7%	78.7%	84.7%	85.3%	86.7%	88.3%	65-69	נקבה
89.0%	87.8%	88.1%	90.2%	70.8%	74.3%	76.0%	78.5%	84.2%	84.1%	81.3%	83.6%	88.6%	88.4%	89.0%	88.7%	70-74	
87.1%	84.8%	84.6%	87.4%	70.5%	74.3%	77.2%	79.4%	80.4%	75.2%	71.9%	74.9%	88.3%	87.6%	87.8%	88.0%	75-79	
79.0%	77.5%	78.7%	81.2%	68.9%	69.9%	69.2%	77.4%	66.5%	64.3%	62.7%	65.0%	84.1%	85.4%	83.2%	83.8%	80-84	

שיעור ההימצאות של השמנת יתר חולנית בקרב חולי סוכרת בני 65-84

35.2%	40.4%	40.9%	40.8%	32.1%	38.8%	42.1%	42.9%	33.8%	35.5%	36.6%	37.9%	32.1%	32.9%	34.6%	37.3%	65-69	זכר
30.5%	37.7%	42.0%	43.5%	30.4%	35.7%	40.3%	43.3%	29.9%	36.5%	36.6%	38.4%	28.6%	31.5%	33.1%	35.6%	70-74	
29.5%	32.3%	39.1%	39.5%	26.6%	30.9%	36.5%	34.8%	30.3%	26.9%	31.2%	36.2%	24.1%	27.3%	31.3%	33.3%	75-79	
26.9%	30.4%	31.2%	31.7%	21.6%	27.0%	32.3%	34.2%	16.2%	25.8%	31.0%	38.4%	20.9%	23.8%	27.6%	29.2%	80-84	
41.6%	48.7%	57.3%	64.0%	39.3%	48.7%	56.6%	59.4%	36.3%	45.7%	56.1%	60.8%	39.1%	43.8%	51.3%	60.7%	65-69	נקבה
36.7%	47.0%	53.8%	61.8%	38.4%	47.3%	54.0%	60.5%	37.0%	43.3%	54.5%	58.0%	37.0%	41.7%	49.7%	57.4%	70-74	
33.1%	47.4%	48.7%	56.0%	35.7%	42.9%	49.8%	58.2%	35.0%	40.3%	48.2%	55.6%	33.8%	39.6%	45.9%	55.1%	75-79	
29.1%	40.9%	45.1%	53.6%	32.6%	40.8%	51.1%	52.8%	27.7%	43.1%	45.8%	49.7%	31.8%	38.4%	43.6%	48.6%	80-84	

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

שיעור קבלת חיסון נגד שפעת בחולי סוכרת בני 18 ומעלה

41.9%	27.0%	25.9%	25.7%	26.1%	19.9%	14.7%	18.0%	-	32.5%	33.7%	40.2%	28.8%	23.3%	21.5%	33.0%	18-34	זכר
45.2%	35.2%	31.9%	29.5%	29.6%	22.6%	17.6%	17.6%	40.0%	40.9%	36.8%	43.2%	30.2%	26.4%	28.6%	37.4%	35-44	
44.0%	34.7%	35.1%	33.4%	32.5%	27.6%	24.5%	25.0%	47.7%	51.2%	50.6%	57.2%	38.8%	35.4%	38.2%	44.3%	45-54	
54.1%	46.5%	42.2%	44.2%	49.4%	38.7%	36.0%	35.6%	68.4%	64.3%	65.0%	65.2%	54.8%	50.8%	49.5%	54.3%	55-64	
75.0%	66.7%	59.9%	57.1%	70.5%	60.2%	52.8%	50.1%	83.0%	76.8%	76.2%	77.6%	75.6%	69.9%	64.3%	66.2%	65-74	
82.5%	72.0%	67.0%	61.5%	78.3%	68.6%	59.9%	56.7%	87.9%	82.4%	78.3%	80.7%	83.8%	77.1%	72.6%	71.9%	75-84	
79.3%	72.4%	66.9%	67.6%	78.9%	68.2%	61.4%	55.2%	90.4%	83.6%	80.1%	73.1%	84.1%	78.8%	74.3%	72.6%	85+	
43.0%	35.2%	26.3%	34.9%	30.7%	22.5%	18.3%	19.3%	71.4%	48.9%	46.1%	55.1%	34.0%	24.3%	27.2%	37.0%	18-34	נקבה
39.8%	31.6%	28.3%	28.4%	31.7%	24.8%	14.7%	19.2%	56.1%	57.1%	52.8%	52.1%	38.6%	30.3%	29.9%	38.2%	35-44	
46.0%	33.8%	31.7%	34.7%	33.8%	26.4%	22.9%	24.1%	60.4%	55.8%	61.9%	64.9%	42.4%	36.8%	38.7%	45.7%	45-54	
52.5%	44.6%	41.2%	41.0%	48.1%	37.2%	32.6%	31.5%	70.8%	69.0%	68.0%	71.9%	54.6%	47.9%	46.5%	52.8%	55-64	
71.5%	62.0%	54.2%	49.1%	68.8%	56.3%	47.1%	44.3%	83.8%	78.5%	75.6%	76.0%	73.4%	64.2%	58.2%	60.0%	65-74	
78.7%	66.9%	56.8%	55.7%	71.4%	59.3%	51.3%	49.4%	83.8%	80.6%	73.3%	76.1%	78.2%	70.1%	64.7%	64.7%	75-84	
74.2%	66.2%	55.3%	49.0%	70.3%	54.8%	50.7%	50.7%	75.1%	71.7%	72.3%	77.5%	76.5%	71.8%	67.4%	65.9%	85+	

שיעור קבלת חיסון נגד פניאומוקוק בחולי סוכרת בני 65-74

77.0%	76.1%	73.1%	77.6%	82.0%	79.6%	78.6%	81.1%	95.0%	93.7%	91.8%	93.4%	78.0%	76.9%	77.7%	82.4%	65-69	זכר
90.0%	90.2%	88.9%	91.6%	93.2%	92.1%	92.1%	91.5%	99.2%	98.6%	97.9%	97.4%	91.0%	90.7%	89.3%	92.9%	70-74	
76.3%	69.8%	70.3%	77.6%	79.9%	77.4%	74.8%	77.1%	97.6%	94.4%	92.9%	91.7%	78.0%	73.5%	73.6%	78.8%	65-69	נקבה
88.5%	86.1%	85.0%	86.0%	91.5%	88.7%	88.4%	90.2%	99.1%	99.0%	98.3%	97.3%	88.2%	86.3%	86.4%	90.2%	70-74	

שיעור ההימצאות של סוכרת בקרב בני 17-2

0.0%	0.0%	0.1%	0.0%	0.1%	0.1%	0.0%	0.1%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%	0.1%	0.1%	2-9	זכר
0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.3%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	10-17	
0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.1%	2-9	נקבה
0.3%	0.2%	0.3%	0.2%	0.2%	0.2%	0.2%	0.2%	0.3%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	10-17	

שיעור המבקרים במרפאת סוכרת ילדים בקרב חולי סוכרת בני 17-2

-	-	-	-	-	93.3%	-	-	-	-	-	-	-	89.7%	96.4%	78.6%	2-9	זכר
-	70.2%	55.9%	43.1%	96.5%	94.0%	88.7%	87.0%	-	-	77.4%	88.6%	93.3%	91.1%	87.4%	80.5%	10-17	
-	-	-	-	-	-	-	-	-	-	-	-	100.0%	100.0%	78.6%	87.5%	2-9	נקבה
-	68.8%	44.6%	54.9%	94.2%	90.7%	90.2%	83.9%	-	-	83.7%	78.1%	91.7%	89.1%	87.9%	81.5%	10-17	

מאחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

מחלות זיהומיות

היקף השימוש בתרופות אנטיביוטיות סיסטמיות, DDD ל-1000 איש, ליום

8.5	9.7	9.8	8.6	9.6	10	10.1	10	9.1	10	9.8	9.6	9	9.9	10	9.4	20-29
9.4	10.7	11	11.2	11.2	11.8	11.9	12.7	10.9	11.2	11	12.1	10.3	11.2	12.1	13.2	30-39
9.1	10.7	11.8	12.9	10.7	11.9	12.3	14.3	10.2	12.3	12.9	14.7	10.2	12	13.6	16.1	40-49
10.5	13.3	15.3	17.1	12.8	14.1	15.3	17.7	12.2	14.8	16.6	19.7	13.6	15	17.7	20.8	50-59
16.8	18.1	20.9	23.8	18	19.1	19.7	23.5	19.1	19.9	21.7	25.8	19.8	20.7	22.4	26.6	60-69
21.3	21.8	26.4	33	23	23.5	22.4	27.2	21.9	23.7	24.5	31.7	24.9	25.4	27	33.4	70-79
23	28.7	30.9	31.8	25.5	24.5	24.9	26.4	24.7	26	28	29.6	29.8	28.8	29.8	38.1	80-84
27.3	29.3	41.1	45.7	28.1	24.6	25.5	31.4	26.6	25.2	28.2	31.4	32.3	30.7	32.5	34.9	85+
17.3	18.6	18.5	16	17.8	18.3	18.9	17.1	15.6	17.6	17	17.4	17.9	19.3	18.6	16.9	20-29
17.4	19.8	21.2	18.4	18.8	19.7	19.4	18.7	19.9	19.8	19.4	20.7	18.9	20.6	21.5	21.4	30-39
15.9	19.2	20.4	19	17.6	19.2	20	19.9	16.4	18.6	21.1	23.3	18.1	20.9	22.9	24.4	40-49
17.9	21.2	23.8	23.4	19.4	22	22	23.7	20.3	22.7	24.5	26.5	23	25.9	27.2	28.2	50-59
25.6	26.3	27.3	30.1	26.1	25.7	24.4	24.9	26.2	26	26.7	30.5	30.5	30.9	30.9	34.7	60-69
31.1	30.6	30.7	36.2	29.5	26.7	24.8	25.7	24.5	25.8	26.4	28.1	36.9	35	33.5	38	70-79
34.8	30.9	36.7	40.5	30.6	28.1	26.3	25.9	23.1	24.7	27	28.9	41.8	39.8	37.3	38.6	80-84
38.8	40	42.8	42.2	31.5	28.9	26.2	32.1	25.3	24	29.5	28.3	43.9	42.2	39.1	38	85+

שיעור השימוש בתרופות אנטיביוטיות מסוג צפלוספורינים וקווינלונים, מתוך סך התרופות האנטיביוטיות

5.5%	4.8%	5.9%	5.5%	19.5%	17.6%	21.0%	14.2%	7.3%	6.5%	6.7%	6.2%	8.6%	6.9%	6.1%	4.6%	0-9
11.4%	13.3%	14.3%	14.6%	13.0%	12.6%	14.2%	14.1%	11.0%	9.5%	13.8%	13.9%	13.2%	12.2%	12.5%	14.0%	10-19
23.0%	20.7%	21.1%	21.2%	18.1%	16.1%	14.8%	15.9%	15.6%	16.5%	16.0%	16.8%	20.1%	17.6%	16.1%	18.0%	20-29
24.3%	23.3%	24.7%	24.8%	17.2%	16.4%	15.7%	17.2%	20.2%	18.8%	17.5%	19.9%	18.6%	18.3%	17.4%	18.9%	30-39
29.7%	29.4%	30.1%	29.6%	20.9%	19.0%	18.9%	20.1%	25.9%	21.1%	22.7%	24.4%	20.8%	21.3%	20.9%	22.2%	40-49
32.5%	34.4%	37.4%	34.7%	22.9%	22.7%	22.7%	23.7%	23.1%	28.9%	25.9%	27.3%	24.9%	25.7%	25.3%	26.5%	50-59
38.6%	40.3%	40.2%	39.4%	28.0%	28.1%	29.1%	29.9%	32.9%	31.8%	33.2%	29.4%	29.6%	30.1%	31.8%	31.8%	60-69
42.0%	44.8%	47.0%	48.5%	32.5%	34.5%	34.2%	32.8%	39.4%	35.1%	39.5%	39.6%	33.1%	35.3%	37.1%	36.9%	70-79
47.6%	45.3%	47.3%	52.2%	36.1%	40.1%	37.6%	40.2%	39.1%	44.9%	38.9%	35.5%	39.4%	42.8%	43.3%	44.5%	80-84
47.9%	51.5%	48.8%	49.7%	42.9%	45.4%	45.6%	46.1%	36.8%	49.5%	52.8%	42.4%	42.9%	47.9%	48.6%	47.3%	85+
8.1%	7.5%	8.1%	8.3%	22.0%	19.8%	17.5%	18.5%	10.2%	9.5%	8.5%	8.4%	11.7%	10.0%	9.1%	8.1%	0-9
16.4%	15.7%	15.2%	15.7%	17.4%	16.3%	15.7%	13.5%	15.3%	14.5%	13.3%	14.1%	16.9%	15.8%	15.4%	15.9%	10-19
33.2%	32.8%	32.2%	31.8%	22.3%	21.6%	21.2%	22.6%	27.2%	26.2%	25.4%	28.2%	25.2%	24.7%	24.3%	26.2%	20-29
33.5%	34.4%	33.5%	34.0%	21.6%	21.4%	21.9%	23.4%	29.8%	26.7%	27.2%	28.3%	24.5%	25.4%	25.4%	26.1%	30-39
38.9%	37.6%	36.0%	35.1%	24.1%	23.2%	23.1%	24.0%	33.0%	29.2%	28.5%	26.9%	27.4%	27.6%	26.3%	26.8%	40-49
38.0%	38.9%	38.5%	35.6%	26.3%	24.8%	25.4%	25.5%	35.7%	31.8%	30.9%	31.3%	29.1%	29.2%	28.8%	27.8%	50-59
37.7%	40.3%	40.8%	39.3%	28.5%	28.1%	29.3%	28.8%	35.5%	36.1%	35.0%	34.7%	30.5%	31.2%	31.5%	31.2%	60-69
37.6%	39.6%	42.9%	42.5%	31.3%	32.0%	33.1%	34.5%	42.9%	39.2%	40.9%	38.7%	30.0%	32.1%	36.3%	36.4%	70-79
34.3%	41.2%	41.8%	44.7%	34.8%	37.4%	36.6%	38.2%	53.3%	49.2%	46.7%	39.1%	32.8%	34.9%	38.7%	42.0%	80-84
38.8%	42.6%	46.5%	51.9%	41.6%	41.7%	44.4%	42.0%	56.4%	50.3%	47.6%	53.8%	34.1%	38.7%	42.2%	43.0%	85+

שיעור המבוטחים בעלי בדיקת RNA חיובית להפטיטיס C אשר קיבלו טיפול אנטי-ויראלי בחמש השנים האחרונות

72.2%	71.2%	73.7%	73.9%	88.6%	84.0%	83.0%	79.8%	-	74.5%	71.0%	68.5%	78.1%	81.4%	74.3%	69.6%	20-59
-	67.4%	64.8%	-	81.3%	72.6%	76.8%	84.4%	-	64.3%	73.3%	62.9%	79.7%	77.9%	72.3%	71.1%	60+
-	79.8%	76.1%	76.2%	89.4%	89.2%	90.3%	85.0%	-	86.2%	71.3%	63.9%	85.9%	87.3%	80.0%	76.1%	20-59
-	76.2%	59.1%	-	75.9%	85.3%	82.8%	81.5%	-	70.8%	76.0%	70.5%	81.8%	80.5%	74.7%	69.6%	60+

היקף השימוש בתרופות אנטיביוטיות סיסטמיות, רכישות ל-1000 איש לשנה

874	967	866	807	848	883	813	818	1090	1069	925	880	790	887	891	931	0-9
352	412	418	383	332	329	329	373	355	420	391	401	306	331	337	350	10-19
367	423	433	413	379	391	399	418	354	400	400	414	341	381	390	385	20-29
406	475	502	551	447	464	465	516	427	448	460	528	402	442	476	543	30-39
397	464	539	650	431	465	476	588	420	493	551	653	399	465	538	663	40-49
460	580	694	867	502	554	594	710	483	603	705	846	510	574	690	837	50-59
731	763	904	1137	722	743	758	909	780	828	907	1098	743	779	855	1057	60-69
909	928	1100	1485	887	901	853	1047	941	987	1026	1357	913	948	1009	1320	70-79
999	1159	1214	1377	1003	949	952	1023	1021	1131	1213	1150	1065	1063	1115	1436	80-84
1098	1152	1468	1789	1082	992	1003	1167	1079	1137	1198	1419	1139	1109	1161	1302	85+
867	997	879	806	873	895	832	796	1039	1094	941	846	804	904	893	883	0-9
442	493	486	444	410	428	423	409	459	527	477	476	378	427	410	392	10-19
762	833	813	723	791	804	811	736	723	807	762	786	713	780	754	681	20-29
796	902	942	867	832	861	829	821	894	878	861	935	772	834	869	875	30-39
723	868	915	927	769	834	839	852	776	849	934	1049	718	822	901	987	40-49
820	974	1073	1167	854	943	914	1003	930	1036	1088	1205	886	998	1063	1143	50-59
1121	1132	1183	1412	1121	1089	1010	1045	1224	1190	1184	1412	1134	1153	1182	1378	60-69
1245	1231	1236	1496	1236	1112	1014	1050	1256	1243	1247	1325	1302	1248	1234	1460	70-79
1316	1207	1372	1650	1245	1140	1071	1013	1205	1246	1289	1274	1394	1357	1314	1421	80-84
1454	1471	1580	1616	1282	1176	1082	1267	1285	1272	1359	1467	1425	1416	1366	1422	85+

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

פרופרציית השימוש בתרופות אנטיביוטיות למיקוד (Antibiotics of concern)

34.8%	33.6%	28.4%	32.2%	28.5%	28.1%	28.1%	28.2%	30.0%	30.3%	27.4%	29.3%	24.9%	26.2%	25.6%	30.6%	0-9
36.4%	36.3%	32.1%	38.4%	31.6%	31.8%	29.8%	34.1%	28.9%	31.0%	30.5%	33.2%	24.8%	25.3%	27.2%	36.0%	10-19
54.1%	53.0%	52.1%	57.2%	50.1%	47.4%	46.0%	48.6%	46.4%	51.3%	49.1%	50.3%	42.7%	43.7%	44.6%	52.0%	20-29
54.4%	54.9%	55.4%	60.8%	46.6%	47.4%	48.5%	52.7%	52.5%	54.9%	52.0%	57.9%	41.6%	44.3%	46.4%	54.2%	30-39
57.0%	58.1%	60.0%	62.9%	51.0%	53.0%	53.7%	56.4%	55.5%	58.8%	57.2%	57.7%	45.3%	47.5%	50.4%	55.8%	40-49
59.8%	61.6%	61.7%	65.3%	54.3%	55.8%	56.6%	58.6%	55.1%	59.2%	56.1%	60.4%	49.2%	50.7%	52.4%	56.5%	50-59
60.5%	63.3%	61.7%	65.1%	56.9%	57.3%	59.0%	57.4%	56.2%	61.2%	60.4%	60.1%	49.2%	50.6%	51.7%	54.3%	60-69
60.0%	58.9%	61.5%	62.1%	57.7%	58.1%	57.4%	57.5%	58.9%	59.9%	59.4%	57.1%	50.0%	50.6%	51.2%	52.7%	70-79
60.3%	60.2%	59.2%	64.4%	56.4%	57.9%	56.4%	57.0%	58.4%	61.8%	57.0%	58.5%	49.0%	51.5%	51.6%	53.4%	80-84
59.4%	57.0%	61.1%	61.0%	59.3%	60.0%	59.1%	57.4%	56.7%	59.7%	65.9%	57.3%	49.7%	53.2%	54.2%	55.8%	85+
32.4%	31.4%	26.2%	29.4%	25.3%	26.0%	25.9%	26.4%	24.7%	27.7%	25.9%	26.6%	22.6%	24.0%	23.8%	28.5%	0-9
35.3%	34.2%	29.5%	35.1%	27.5%	28.5%	27.3%	30.7%	30.6%	30.8%	29.0%	31.7%	23.1%	25.0%	26.1%	34.3%	10-19
51.4%	51.0%	48.0%	51.9%	40.8%	40.2%	39.6%	42.1%	47.9%	48.7%	46.4%	47.3%	38.0%	39.2%	40.0%	47.2%	20-29
51.8%	53.8%	53.0%	56.7%	40.8%	42.5%	43.6%	48.0%	49.3%	50.6%	50.6%	51.9%	38.9%	41.7%	42.8%	48.7%	30-39
55.1%	57.6%	56.7%	61.2%	42.6%	45.6%	48.4%	51.7%	51.7%	55.3%	54.9%	56.9%	41.4%	44.6%	46.0%	51.1%	40-49
55.4%	58.6%	59.6%	63.1%	45.9%	48.7%	51.0%	53.0%	55.6%	56.2%	56.0%	57.3%	43.7%	46.5%	47.4%	52.0%	50-59
57.0%	57.5%	59.2%	62.0%	48.6%	50.1%	51.8%	51.8%	53.3%	57.0%	56.5%	56.2%	44.0%	45.5%	46.0%	48.5%	60-69
51.3%	55.1%	56.8%	59.0%	47.7%	48.4%	50.1%	49.9%	51.5%	56.1%	53.0%	53.6%	41.2%	43.1%	44.5%	46.9%	70-79
49.9%	52.2%	51.7%	55.1%	46.3%	47.7%	49.7%	49.4%	53.0%	53.8%	54.4%	54.2%	40.6%	42.3%	44.5%	46.0%	80-84
49.8%	51.3%	52.4%	59.5%	48.8%	49.4%	50.6%	52.7%	53.7%	50.6%	53.5%	51.9%	41.0%	43.5%	46.5%	48.5%	85+

זכר

נקבה

מאוחדת				מכבי				לאומית				כללית				SES גיל
4	3	2	1	4	3	2	1	4	3	2	1	4	3	2	1	

בריאות הנפש

שיעור ההימצאות של מחלת נפש קשה (SMI) בקרב בני 18 ומעלה

0.5%	0.6%	0.7%	0.6%	0.8%	1.1%	1.6%	1.7%	1.8%	1.9%	2.1%	2.9%	0.7%	1.2%	1.1%	0.7%	18-24
0.5%	0.8%	1.0%	1.1%	1.2%	1.5%	2.2%	2.5%	2.4%	2.5%	3.0%	3.2%	1.2%	1.7%	1.8%	1.5%	25-34
1.0%	1.3%	1.6%	1.3%	1.3%	2.0%	2.9%	3.3%	2.2%	2.8%	3.4%	3.7%	1.3%	1.9%	2.5%	2.1%	35-44
0.8%	1.2%	1.7%	1.5%	1.2%	1.9%	3.0%	3.4%	2.1%	2.6%	3.2%	3.5%	1.4%	2.2%	2.9%	2.3%	45-54
0.6%	0.7%	1.3%	1.5%	1.1%	1.6%	2.2%	2.7%	1.6%	1.8%	3.0%	3.8%	1.4%	2.3%	2.8%	2.5%	55-64
0.5%	0.6%	0.9%	0.8%	0.9%	1.2%	1.4%	1.6%	1.2%	1.3%	1.6%	2.1%	0.9%	1.2%	1.6%	1.5%	65+
0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	0.8%	0.4%	0.9%	1.2%	0.8%	0.7%	0.5%	0.6%	0.5%	0.3%	18-24
0.4%	0.5%	0.4%	0.4%	0.8%	1.1%	1.2%	1.1%	1.1%	1.7%	1.6%	1.4%	0.7%	0.9%	0.9%	0.7%	25-34
0.7%	0.7%	0.7%	0.7%	1.0%	1.4%	1.9%	1.9%	1.2%	1.5%	2.0%	2.3%	0.8%	1.2%	1.4%	1.2%	35-44
0.6%	0.8%	1.2%	1.1%	1.1%	1.6%	2.3%	2.4%	1.8%	2.0%	2.4%	2.3%	1.0%	1.6%	2.1%	1.6%	45-54
0.7%	0.9%	1.2%	1.0%	1.3%	1.7%	2.2%	2.7%	1.5%	1.9%	2.4%	3.0%	1.3%	1.9%	2.3%	2.0%	55-64
0.8%	0.8%	0.9%	0.9%	1.2%	1.5%	1.7%	1.8%	1.5%	1.6%	1.8%	2.3%	1.1%	1.4%	1.7%	1.5%	65+

שיעור ההימצאות של סוכרת בקרב חולי SMI, בני 18 ומעלה

3.2%	1.1%	1.1%	2.9%	2.1%	1.9%	1.9%	2.7%	0.9%	0.3%	0.5%	1.7%	1.8%	1.8%	2.1%	2.2%	18-34
5.1%	6.4%	7.1%	8.8%	4.2%	5.0%	6.9%	5.0%	2.6%	3.2%	5.9%	5.3%	7.0%	5.9%	9.0%	9.1%	35-44
10.8%	16.3%	16.7%	13.8%	10.2%	13.7%	17.0%	15.9%	6.4%	14.7%	17.6%	19.1%	13.7%	18.4%	16.8%	20.0%	45-54
10.4%	24.3%	26.5%	31.2%	18.9%	23.0%	25.6%	30.2%	31.0%	23.4%	26.6%	25.7%	21.3%	28.3%	28.2%	31.6%	55-64
33.3%	33.3%	29.7%	35.7%	27.4%	36.1%	29.3%	32.1%	45.5%	43.8%	32.5%	40.4%	37.1%	33.2%	33.4%	31.2%	65+
0.0%	4.5%	3.3%	2.2%	1.7%	1.0%	1.2%	1.8%	2.0%	1.5%	2.8%	1.8%	2.7%	2.5%	3.8%	3.9%	18-34
3.4%	6.8%	7.2%	3.7%	3.0%	3.9%	6.9%	9.6%	2.4%	4.5%	6.6%	9.8%	4.4%	6.6%	9.0%	9.3%	35-44
5.6%	8.2%	16.7%	19.0%	6.4%	10.8%	15.0%	17.2%	13.9%	9.2%	12.7%	15.9%	12.6%	15.1%	18.6%	21.5%	45-54
29.1%	29.5%	23.5%	26.7%	15.3%	19.2%	25.5%	28.2%	25.4%	25.1%	29.3%	26.1%	22.5%	27.1%	31.1%	35.8%	55-64
13.5%	30.9%	29.8%	37.0%	24.1%	28.6%	29.9%	35.0%	29.3%	29.1%	31.6%	37.1%	29.0%	33.6%	38.4%	39.2%	65+

שיעור התייעוד של מרכיבי BMI בקרב חולי SMI, בני 20-64

47.6%	41.1%	47.5%	44.6%	27.2%	27.1%	28.1%	26.8%	32.5%	31.5%	28.5%	25.2%	35.5%	32.8%	33.0%	29.7%	20-34
53.8%	52.4%	51.5%	53.3%	33.0%	32.1%	33.8%	34.2%	30.4%	33.1%	33.5%	29.5%	40.2%	38.0%	41.5%	41.4%	35-44
62.0%	69.9%	61.4%	68.7%	38.1%	43.6%	45.4%	50.5%	40.4%	48.8%	45.8%	48.4%	56.9%	60.8%	60.1%	61.2%	45-64
42.1%	55.5%	67.7%	67.7%	34.8%	38.2%	43.4%	47.6%	29.8%	41.4%	42.3%	48.8%	40.1%	45.5%	46.2%	42.4%	20-34
60.0%	63.0%	63.4%	74.3%	38.3%	39.0%	48.2%	50.6%	40.5%	41.0%	44.8%	49.5%	46.0%	50.8%	51.8%	48.6%	35-44
71.3%	68.0%	69.9%	74.5%	44.5%	46.4%	53.1%	55.7%	53.4%	52.8%	56.1%	54.9%	63.7%	67.7%	68.7%	67.4%	45-64

שיעור ההימצאות של השמנת יתר בקרב חולי SMI, בני 20-64

26.7%	35.9%	36.9%	35.3%	38.0%	41.6%	37.7%	39.3%	18.9%	39.0%	31.3%	37.0%	35.7%	35.6%	37.5%	38.9%	20-34
27.9%	45.4%	48.7%	52.7%	37.8%	46.7%	44.6%	50.7%	-	39.3%	38.9%	41.5%	36.2%	38.0%	40.0%	37.6%	35-44
35.2%	44.3%	43.3%	45.6%	40.0%	40.2%	47.7%	46.3%	32.7%	39.8%	40.3%	44.0%	37.0%	35.5%	36.1%	37.5%	45-64
-	39.3%	59.5%	52.2%	33.3%	38.5%	40.5%	49.2%	-	41.7%	52.3%	40.0%	36.6%	41.5%	47.2%	50.0%	20-34
30.6%	46.7%	48.9%	51.9%	40.7%	48.2%	51.9%	49.6%	-	45.5%	51.3%	66.7%	42.5%	45.3%	51.6%	58.4%	35-44
33.7%	51.3%	53.1%	64.2%	37.6%	49.6%	52.0%	62.0%	37.1%	40.0%	54.1%	58.5%	43.3%	47.9%	51.3%	58.6%	45-64

שיעור המפגשים הטיפוליים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך

-	26.1%	37.5%	27.0%	35.4%	42.2%	40.7%	26.3%	-	16.7%	17.2%	14.9%	68.8%	76.9%	77.8%	81.9%	18-34
-	31.6%	38.9%	26.5%	36.1%	46.1%	33.6%	37.1%	-	14.0%	12.2%	13.2%	71.3%	76.7%	78.8%	77.1%	35-54
-	-	-	-	40.0%	28.3%	33.3%	-	-	-	-	-	67.2%	71.1%	71.4%	81.3%	55+
-	40.0%	32.7%	43.6%	45.4%	46.2%	39.7%	37.3%	-	-	20.8%	8.3%	78.1%	77.9%	75.5%	78.9%	18-34
-	37.0%	37.0%	-	51.0%	37.2%	35.4%	32.6%	-	-	24.4%	-	76.0%	77.3%	81.9%	77.8%	35-54
-	-	26.7%	-	37.5%	37.5%	30.2%	-	-	-	6.5%	-	76.4%	69.1%	68.6%	73.5%	55+